



**Multicomponent diastereoselective synthesis of  
indolizidines via 1,3-dipolar cycloadditions of azomethine  
ylides**

Journal:	<i>SYNTHESIS</i>
Manuscript ID	SS-2016-07-0541-OP.R1
Manuscript Type:	Original Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Selva, Veronica; Universitat d'Alacant Facultat de Ciències, Organic Chemistry Castello, Luis; University of Alicante, Organic Chemistry Najera, Carmen; University of Alicante, Organic Chemistry Sansano, Jose; University of Alicante, Organic Chemistry
Keywords:	cycloaddition, fused-ring systems, heterocycles, multicomponent reaction, diastereoselectivity
Abstract:	The synthesis of polyfunctionalized indolizidines from pipecolinic acid alkyl ester derivatives, aldehydes and a wide range of dipolarophiles by a multicomponent 1,3-dipolar cycloadditions has been developed in a diastereoselective manner. Reactions take place in toluene with short reaction times at 70 °C, giving good yields. The synthesis of these fused heterocycles is also studied starting from the pipecolinic acid, generating the dipole through a decarboxylative route at 120 °C. The relative configuration of the resulting products, as well as the mechanistic pathways are also explained.

SCHOLARONE™  
Manuscripts


## SUPPORTING INFORMATION


Multicomponent diastereoselective synthesis of indolizidines via  
1,3-dipolar cycloadditions of azomethine ylidesCastelló, Luis M.<sup>a,b</sup> Selva, Verónica<sup>a,b</sup> Nájera, Carmen<sup>a</sup> \* Sansano, José M.<sup>a,b</sup><sup>a</sup> *Departamento de Química Orgánica. Facultad de Ciencias, Universidad de Alicante, 03080-Alicante, Spain. Centro de Innovación en Química Avanzada (ORFEO-CINQA).*<sup>b</sup> *Instituto de Síntesis Orgánica (ISO). Universidad de Alicante, 03080-Alicante, Spain.*


\* cnajera@ua.es.


## Table of contents


1. <sup>1</sup> H and <sup>13</sup> C NMR spectra .....	S2
2. X-Ray diffraction analysis of compound <i>endo</i> -13 .....	S25


1.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra





































































## 2. X-Ray diffraction analysis of compound *endo*-13



```

data_x

_audit_creation_method          SHELXL-97
_chemical_name_systematic
;
?
;
_chemical_name_common           ?
_chemical_melting_point         ?
_chemical_formula_moiety
'C18 H22 N2 O5'
_chemical_formula_sum
'C18 H22 N2 O5'
_chemical_formula_weight        346.38

loop_
_atom_type_symbol
_atom_type_description
_atom_type_scatter_dispersion_real
_atom_type_scatter_dispersion_imag
_atom_type_scatter_source
'C' 'C' 0.0033 0.0016
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'H' 'H' 0.0000 0.0000
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'N' 'N' 0.0061 0.0033
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'O' 'O' 0.0106 0.0060
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

_symmetry_cell_setting          'triclinic'
_symmetry_space_group_name_H-M 'P -1'

loop_
_symmetry_equiv_pos_as_xyz
'x, y, z'
'-x, -y, -z'

_cell_length_a                  8.627(3)
_cell_length_b                  10.120(4)
_cell_length_c                  10.797(4)
_cell_angle_alpha               87.973(7)
_cell_angle_beta               86.925(7)
_cell_angle_gamma               67.294(6)
_cell_volume                    868.2(6)
_cell_formula_units_Z           2
_cell_measurement_temperature    298(1)
_cell_measurement_reflns_used    1391
_cell_measurement_theta_min      2.56
_cell_measurement_theta_max      24.24

_exptl_crystal_description      plate
_exptl_crystal_colour           colorless
_exptl_crystal_size_max         0.61
_exptl_crystal_size_mid         0.33
_exptl_crystal_size_min         0.11
_exptl_crystal_density_meas     ?
_exptl_crystal_density_diffrn   1.325
_exptl_crystal_density_method   'not measured'

```

```

_exptl_crystal_F_000          368
_exptl_absorpt_coefficient_mu 0.097
_exptl_absorpt_correction_type 'multi-scan'
_exptl_absorpt_correction_T_min 0.8331
_exptl_absorpt_correction_T_max 0.9894
_exptl_absorpt_process_details 'Bruker/Siemens SADABS V2.03'

_exptl_special_details
;
?
;

_diffrn_ambient_temperature    298(1)
_diffrn_radiation_wavelength    0.71073
_diffrn_radiation_type          MoK\alpha
_diffrn_radiation_source        'fine-focus sealed tube'
_diffrn_radiation_monochromator graphite
_diffrn_measurement_device_type 'CCD area detector'
_diffrn_measurement_method      'phi and omega scans'
_diffrn_detector_area_resol_mean ?
_diffrn_standards_number        ?
_diffrn_standards_interval_count ?
_diffrn_standards_interval_time ?
_diffrn_standards_decay_%       ?
_diffrn_reflns_number           6059
_diffrn_reflns_av_R_equivalents 0.0264
_diffrn_reflns_av_sigmaI/netI   0.0422
_diffrn_reflns_limit_h_min      -10
_diffrn_reflns_limit_h_max      10
_diffrn_reflns_limit_k_min      -12
_diffrn_reflns_limit_k_max      12
_diffrn_reflns_limit_l_min      -12
_diffrn_reflns_limit_l_max      12
_diffrn_reflns_theta_min        1.89
_diffrn_reflns_theta_max        25.06
_reflns_number_total            3060
_reflns_number_gt               2127
_reflns_threshold_expression     'I>2\sigma(I)'

_computing_data_collection      'Bruker SMART V5.625'
_computing_cell_refinement      'Bruker SMART V5.625'
_computing_data_reduction       'Bruker SAINT V6.28A'
_computing_structure_solution   'SHELXS-97 (Sheldrick, 1997)'
_computing_structure_refinement 'SHELXL-97 (Sheldrick, 1997)'
_computing_molecular_graphics   'SHELXTL Rel. 6.12/V (Siemens, 1996)'
_computing_publication_material 'SHELX-97 (Sheldrick, 1997)'

_refine_special_details
;
Refinement of F2 against ALL reflections. The weighted R-factor wR and
goodness of fit S are based on F2, conventional R-factors R are based
on F, with F set to zero for negative F2. The threshold expression of
F2 > 2sigma(F2) is used only for calculating R-factors(gt) etc. and is
not relevant to the choice of reflections for refinement. R-factors based
on F2 are statistically about twice as large as those based on F, and R-
factors based on ALL data will be even larger.
;

_refine_ls_structure_factor_coef Fsqd
_refine_ls_matrix_type          full

```

```

_refine_ls_weighting_scheme      calc
_refine_ls_weighting_details
'calc w=1/[\s^2^(Fo^2^)+(0.0756P)^2^] where P=(Fo^2^+2Fc^2^)/3'
_atom_sites_solution_primary    direct
_atom_sites_solution_secondary  difmap
_atom_sites_solution_hydrogens  geom
_refine_ls_hydrogen_treatment   constr
_refine_ls_extinction_method     none
_refine_ls_extinction_coef       ?
_refine_ls_number_reflns        3060
_refine_ls_number_parameters     250
_refine_ls_number_restraints     0
_refine_ls_R_factor_all         0.0742
_refine_ls_R_factor_gt          0.0500
_refine_ls_wR_factor_ref        0.1417
_refine_ls_wR_factor_gt         0.1274
_refine_ls_goodness_of_fit_ref   1.043
_refine_ls_restrained_S_all     1.043
_refine_ls_shift/su_max         0.012
_refine_ls_shift/su_mean        0.001

```

```

loop_
_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
_atom_site_adp_type
_atom_site_occupancy
_atom_site_symmetry_multiplicity
_atom_site_calc_flag
_atom_site_refinement_flags
_atom_site_disorder_assembly
_atom_site_disorder_group
C1 C 1.1659(3) 0.1139(2) 0.0746(2) 0.0416(5) Uani 1 1 d . . .
O1 O 1.2020(2) 0.06866(18) -0.02970(15) 0.0595(5) Uani 1 1 d . . .
N1 N 1.2536(2) 0.1796(2) 0.13498(17) 0.0455(5) Uani 1 1 d . . .
C2 C 1.2027(3) 0.2052(2) 0.2584(2) 0.0449(5) Uani 1 1 d . . .
O2 O 1.2684(2) 0.2551(2) 0.32914(16) 0.0655(5) Uani 1 1 d . . .
C3 C 1.0621(3) 0.1523(2) 0.28651(19) 0.0399(5) Uani 1 1 d . . .
H3 H 1.1001 0.0682 0.3421 0.048 Uiso 1 1 calc R . .
C4 C 0.8493(3) 0.2265(2) 0.12686(17) 0.0351(5) Uani 1 1 d . . .
H4 H 0.7647 0.1839 0.1353 0.042 Uiso 1 1 calc R . .
C5 C 0.8415(3) 0.2946(2) 0.00120(18) 0.0390(5) Uani 1 1 d . . .
C6 C 0.7682(3) 0.2852(3) -0.1022(2) 0.0589(7) Uani 1 1 d . . .
H6 H 0.7003 0.2344 -0.1131 0.071 Uiso 1 1 calc R . .
C7 C 0.8178(4) 0.3720(3) -0.1944(2) 0.0709(9) Uani 1 1 d . . .
H7 H 0.7855 0.3893 -0.2761 0.085 Uiso 1 1 calc R . .
C8 C 0.9166(4) 0.4211(3) -0.1400(2) 0.0720(8) Uani 1 1 d . . .
H8 H 0.9674 0.4784 -0.1789 0.086 Uiso 1 1 calc R . .
O3 O 0.9349(2) 0.37721(18) -0.01952(14) 0.0584(5) Uani 1 1 d . . .
N2 N 0.8160(2) 0.33295(17) 0.22380(14) 0.0349(4) Uani 1 1 d . . .
C9 C 0.6444(3) 0.4369(2) 0.2394(2) 0.0461(6) Uani 1 1 d . . .
H9A H 0.5733 0.3912 0.2779 0.055 Uiso 1 1 calc R . .
H9B H 0.6009 0.4755 0.1592 0.055 Uiso 1 1 calc R . .
C10 C 0.6429(3) 0.5571(3) 0.3207(2) 0.0537(6) Uani 1 1 d . . .
H10A H 0.7022 0.6102 0.2770 0.064 Uiso 1 1 calc R . .
H10B H 0.5277 0.6226 0.3376 0.064 Uiso 1 1 calc R . .
C11 C 0.7257(3) 0.4985(3) 0.4423(2) 0.0554(6) Uani 1 1 d . . .

```

H11A H 0.6570 0.4578 0.4914 0.066 Uiso 1 1 calc R . .  
 H11B H 0.7326 0.5762 0.4888 0.066 Uiso 1 1 calc R . .  
 C12 C 0.9018(3) 0.3840(2) 0.42078(19) 0.0461(6) Uani 1 1 d . . .  
 H12A H 0.9475 0.3427 0.4998 0.055 Uiso 1 1 calc R . .  
 H12B H 0.9749 0.4271 0.3813 0.055 Uiso 1 1 calc R . .  
 C13 C 0.8956(3) 0.2660(2) 0.33833(17) 0.0366(5) Uani 1 1 d . . .  
 C14 C 1.0236(3) 0.1111(2) 0.15965(18) 0.0376(5) Uani 1 1 d . . .  
 H14 H 1.0203 0.0154 0.1634 0.045 Uiso 1 1 calc R . .  
 C15 C 0.8081(3) 0.1800(2) 0.4153(2) 0.0461(6) Uani 1 1 d . A .  
 O4 O 0.8085(3) 0.1675(2) 0.52540(16) 0.0915(8) Uani 1 1 d . . .  
 O5 O 0.7422(2) 0.1141(2) 0.34412(14) 0.0616(5) Uani 1 1 d . . .  
 C16A C 0.6630(14) 0.0282(16) 0.4102(12) 0.053(3) Uani 0.50 1 d P A 1  
 H16A H 0.5542 0.0892 0.4458 0.063 Uiso 0.50 1 calc PR A 1  
 H16B H 0.7324 -0.0261 0.4767 0.063 Uiso 0.50 1 calc PR A 1  
 C17A C 0.6425(9) -0.0733(8) 0.3161(8) 0.067(2) Uani 0.50 1 d P A 1  
 H17A H 0.5718 -0.0183 0.2517 0.100 Uiso 0.50 1 calc PR A 1  
 H17B H 0.5920 -0.1334 0.3572 0.100 Uiso 0.50 1 calc PR A 1  
 H17C H 0.7508 -0.1318 0.2805 0.100 Uiso 0.50 1 calc PR A 1  
 C16B C 0.677(2) 0.0070(19) 0.3987(14) 0.096(6) Uani 0.50 1 d P A 2  
 H16C H 0.6924 -0.0003 0.4874 0.115 Uiso 0.50 1 calc PR A 2  
 H16D H 0.5578 0.0403 0.3858 0.115 Uiso 0.50 1 calc PR A 2  
 C17B C 0.7494(15) -0.1134(9) 0.3528(9) 0.109(4) Uani 0.50 1 d P A 2  
 H17D H 0.7406 -0.1059 0.2644 0.163 Uiso 0.50 1 calc PR A 2  
 H17E H 0.6975 -0.1763 0.3861 0.163 Uiso 0.50 1 calc PR A 2  
 H17F H 0.8659 -0.1510 0.3726 0.163 Uiso 0.50 1 calc PR A 2  
 C18 C 1.3891(3) 0.2141(3) 0.0750(3) 0.0680(8) Uani 1 1 d . . .  
 H18A H 1.4062 0.2860 0.1216 0.102 Uiso 1 1 calc R . .  
 H18B H 1.3597 0.2494 -0.0077 0.102 Uiso 1 1 calc R . .  
 H18C H 1.4907 0.1295 0.0718 0.102 Uiso 1 1 calc R . .

loop\_

\_atom\_site\_aniso\_label  
 \_atom\_site\_aniso\_U\_11  
 \_atom\_site\_aniso\_U\_22  
 \_atom\_site\_aniso\_U\_33  
 \_atom\_site\_aniso\_U\_23  
 \_atom\_site\_aniso\_U\_13  
 \_atom\_site\_aniso\_U\_12  
 C1 0.0402(12) 0.0389(12) 0.0415(12) -0.0031(10) -0.0023(10) -0.0103(10)  
 O1 0.0585(11) 0.0675(11) 0.0525(11) -0.0202(9) 0.0104(8) -0.0241(9)  
 N1 0.0366(10) 0.0522(11) 0.0499(11) -0.0040(9) -0.0001(8) -0.0195(9)  
 C2 0.0396(12) 0.0479(13) 0.0444(13) -0.0006(10) -0.0089(10) -0.0129(11)  
 O2 0.0529(11) 0.0883(13) 0.0635(11) -0.0186(10) -0.0118(9) -0.0337(10)  
 C3 0.0425(13) 0.0386(12) 0.0382(12) 0.0068(9) -0.0078(9) -0.0152(10)  
 C4 0.0381(11) 0.0366(11) 0.0340(11) 0.0021(9) -0.0031(9) -0.0183(10)  
 C5 0.0420(12) 0.0413(12) 0.0343(11) -0.0007(9) -0.0008(9) -0.0170(10)  
 C6 0.0518(15) 0.0832(19) 0.0416(13) -0.0060(13) -0.0079(11) -0.0246(14)  
 C7 0.0661(18) 0.097(2) 0.0312(13) 0.0146(14) -0.0079(12) -0.0113(17)  
 C8 0.087(2) 0.0716(18) 0.0468(16) 0.0211(13) 0.0048(15) -0.0218(17)  
 O3 0.0750(12) 0.0603(11) 0.0475(10) 0.0090(8) 0.0013(8) -0.0357(10)  
 N2 0.0389(10) 0.0345(9) 0.0310(9) -0.0003(7) -0.0033(7) -0.0135(8)  
 C9 0.0408(13) 0.0454(13) 0.0468(13) -0.0019(10) -0.0027(10) -0.0106(11)  
 C10 0.0518(15) 0.0459(14) 0.0589(15) -0.0107(11) 0.0083(12) -0.0144(12)  
 C11 0.0600(16) 0.0576(15) 0.0536(14) -0.0176(12) 0.0106(12) -0.0284(13)  
 C12 0.0550(14) 0.0552(14) 0.0362(12) -0.0044(10) -0.0014(10) -0.0298(12)  
 C13 0.0417(12) 0.0426(12) 0.0294(10) 0.0027(9) -0.0043(9) -0.0207(10)  
 C14 0.0418(12) 0.0329(11) 0.0396(12) 0.0017(9) -0.0038(9) -0.0160(10)  
 C15 0.0573(15) 0.0492(13) 0.0366(13) 0.0030(10) -0.0014(10) -0.0262(12)  
 O4 0.160(2) 0.1198(18) 0.0362(11) 0.0124(10) -0.0036(11) -0.1007(18)  
 O5 0.0823(13) 0.0842(13) 0.0450(9) 0.0116(9) -0.0022(9) -0.0626(12)

C16A 0.050(4) 0.058(5) 0.060(6) 0.014(4) -0.017(4) -0.032(4)  
C17A 0.079(5) 0.060(5) 0.073(5) 0.003(4) 0.008(4) -0.042(4)  
C16B 0.166(12) 0.121(12) 0.057(6) 0.001(6) 0.028(6) -0.123(11)  
C17B 0.182(11) 0.050(5) 0.095(7) -0.001(5) 0.045(7) -0.051(7)  
C18 0.0486(16) 0.086(2) 0.0781(19) -0.0125(15) 0.0145(13) -0.0362(15)

\_geom\_special\_details

;

All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

;

loop\_

\_geom\_bond\_atom\_site\_label\_1  
\_geom\_bond\_atom\_site\_label\_2  
\_geom\_bond\_distance  
\_geom\_bond\_site\_symmetry\_2  
\_geom\_bond\_publ\_flag

C1 O1 1.210(2) . ?  
C1 N1 1.382(3) . ?  
C1 C14 1.501(3) . ?  
N1 C2 1.382(3) . ?  
N1 C18 1.458(3) . ?  
C2 O2 1.206(3) . ?  
C2 C3 1.517(3) . ?  
C3 C14 1.535(3) . ?  
C3 C13 1.544(3) . ?  
C3 H3 0.9800 . ?  
C4 N2 1.465(3) . ?  
C4 C5 1.492(3) . ?  
C4 C14 1.556(3) . ?  
C4 H4 0.9800 . ?  
C5 C6 1.336(3) . ?  
C5 O3 1.373(3) . ?  
C6 C7 1.459(4) . ?  
C6 H6 0.9300 . ?  
C7 C8 1.311(4) . ?  
C7 H7 0.9300 . ?  
C8 O3 1.357(3) . ?  
C8 H8 0.9300 . ?  
N2 C9 1.453(3) . ?  
N2 C13 1.460(2) . ?  
C9 C10 1.521(3) . ?  
C9 H9A 0.9700 . ?  
C9 H9B 0.9700 . ?  
C10 C11 1.518(3) . ?  
C10 H10A 0.9700 . ?  
C10 H10B 0.9700 . ?  
C11 C12 1.526(3) . ?  
C11 H11A 0.9700 . ?  
C11 H11B 0.9700 . ?  
C12 C13 1.533(3) . ?  
C12 H12A 0.9700 . ?  
C12 H12B 0.9700 . ?  
C13 C15 1.551(3) . ?  
C14 H14 0.9800 . ?

C15 O4 1.191(3) . ?  
C15 O5 1.317(3) . ?  
O5 C16A 1.446(14) . ?  
O5 C16B 1.490(15) . ?  
C16A C17A 1.534(17) . ?  
C16A H16A 0.9700 . ?  
C16A H16B 0.9700 . ?  
C17A H17A 0.9600 . ?  
C17A H17B 0.9600 . ?  
C17A H17C 0.9600 . ?  
C16B C17B 1.242(19) . ?  
C16B H16C 0.9700 . ?  
C16B H16D 0.9700 . ?  
C17B H17D 0.9600 . ?  
C17B H17E 0.9600 . ?  
C17B H17F 0.9600 . ?  
C18 H18A 0.9600 . ?  
C18 H18B 0.9600 . ?  
C18 H18C 0.9600 . ?

loop\_  
\_geom\_angle\_atom\_site\_label\_1  
\_geom\_angle\_atom\_site\_label\_2  
\_geom\_angle\_atom\_site\_label\_3  
\_geom\_angle  
\_geom\_angle\_site\_symmetry\_1  
\_geom\_angle\_site\_symmetry\_3  
\_geom\_angle\_publ\_flag  
O1 C1 N1 123.7(2) . . ?  
O1 C1 C14 127.9(2) . . ?  
N1 C1 C14 108.42(18) . . ?  
C2 N1 C1 113.06(19) . . ?  
C2 N1 C18 123.7(2) . . ?  
C1 N1 C18 123.2(2) . . ?  
O2 C2 N1 123.7(2) . . ?  
O2 C2 C3 128.1(2) . . ?  
N1 C2 C3 108.08(18) . . ?  
C2 C3 C14 104.40(17) . . ?  
C2 C3 C13 115.15(17) . . ?  
C14 C3 C13 106.01(16) . . ?  
C2 C3 H3 110.3 . . ?  
C14 C3 H3 110.3 . . ?  
C13 C3 H3 110.3 . . ?  
N2 C4 C5 110.70(16) . . ?  
N2 C4 C14 103.07(15) . . ?  
C5 C4 C14 115.56(17) . . ?  
N2 C4 H4 109.1 . . ?  
C5 C4 H4 109.1 . . ?  
C14 C4 H4 109.1 . . ?  
C6 C5 O3 110.6(2) . . ?  
C6 C5 C4 132.7(2) . . ?  
O3 C5 C4 116.48(18) . . ?  
C5 C6 C7 104.9(2) . . ?  
C5 C6 H6 127.5 . . ?  
C7 C6 H6 127.5 . . ?  
C8 C7 C6 106.9(2) . . ?  
C8 C7 H7 126.5 . . ?  
C6 C7 H7 126.5 . . ?  
C7 C8 O3 111.1(3) . . ?  
C7 C8 H8 124.4 . . ?



O3 C8 H8 124.4 . . ?  
C8 O3 C5 106.4(2) . . ?  
C9 N2 C13 115.24(16) . . ?  
C9 N2 C4 116.90(16) . . ?  
C13 N2 C4 111.08(15) . . ?  
N2 C9 C10 108.92(18) . . ?  
N2 C9 H9A 109.9 . . ?  
C10 C9 H9A 109.9 . . ?  
N2 C9 H9B 109.9 . . ?  
C10 C9 H9B 109.9 . . ?  
H9A C9 H9B 108.3 . . ?  
C11 C10 C9 111.1(2) . . ?  
C11 C10 H10A 109.4 . . ?  
C9 C10 H10A 109.4 . . ?  
C11 C10 H10B 109.4 . . ?  
C9 C10 H10B 109.4 . . ?  
H10A C10 H10B 108.0 . . ?  
C10 C11 C12 111.49(19) . . ?  
C10 C11 H11A 109.3 . . ?  
C12 C11 H11A 109.3 . . ?  
C10 C11 H11B 109.3 . . ?  
C12 C11 H11B 109.3 . . ?  
H11A C11 H11B 108.0 . . ?  
C11 C12 C13 110.22(19) . . ?  
C11 C12 H12A 109.6 . . ?  
C13 C12 H12A 109.6 . . ?  
C11 C12 H12B 109.6 . . ?  
C13 C12 H12B 109.6 . . ?  
H12A C12 H12B 108.1 . . ?  
N2 C13 C12 108.17(17) . . ?  
N2 C13 C3 101.03(15) . . ?  
C12 C13 C3 119.06(18) . . ?  
N2 C13 C15 115.88(17) . . ?  
C12 C13 C15 107.92(17) . . ?  
C3 C13 C15 105.11(17) . . ?  
C1 C14 C3 104.49(17) . . ?  
C1 C14 C4 113.67(17) . . ?  
C3 C14 C4 105.92(16) . . ?  
C1 C14 H14 110.8 . . ?  
C3 C14 H14 110.8 . . ?  
C4 C14 H14 110.8 . . ?  
O4 C15 O5 123.5(2) . . ?  
O4 C15 C13 124.4(2) . . ?  
O5 C15 C13 112.05(17) . . ?  
C15 O5 C16A 114.8(6) . . ?  
C15 O5 C16B 120.3(7) . . ?  
C16A O5 C16B 9.1(12) . . ?  
O5 C16A C17A 107.2(9) . . ?  
O5 C16A H16A 110.3 . . ?  
C17A C16A H16A 110.3 . . ?  
O5 C16A H16B 110.3 . . ?  
C17A C16A H16B 110.3 . . ?  
H16A C16A H16B 108.5 . . ?  
C17B C16B O5 112.6(12) . . ?  
C17B C16B H16C 109.1 . . ?  
O5 C16B H16C 109.1 . . ?  
C17B C16B H16D 109.1 . . ?  
O5 C16B H16D 109.1 . . ?  
H16C C16B H16D 107.8 . . ?  
C16B C17B H17D 109.5 . . ?

C16B C17B H17E 109.5 . . ?  
H17D C17B H17E 109.5 . . ?  
C16B C17B H17F 109.5 . . ?  
H17D C17B H17F 109.5 . . ?  
H17E C17B H17F 109.5 . . ?  
N1 C18 H18A 109.5 . . ?  
N1 C18 H18B 109.5 . . ?  
H18A C18 H18B 109.5 . . ?  
N1 C18 H18C 109.5 . . ?  
H18A C18 H18C 109.5 . . ?  
H18B C18 H18C 109.5 . . ?

\_diffn\_measured\_fraction\_theta\_max 0.998  
\_diffn\_refl\_theta\_full 25.00  
\_diffn\_measured\_fraction\_theta\_full 0.999  
\_refine\_diff\_density\_max 0.300  
\_refine\_diff\_density\_min -0.217  
\_refine\_diff\_density\_rms 0.042


# Multicomponent diastereoselective synthesis of indolizidines via 1,3-dipolar cycloadditions of azomethine ylides

Castelló, Luis M.<sup>a,b</sup>  
Selva, Verónica<sup>a,b</sup>  
Nájera, Carmen<sup>a,\*</sup>  
Sansano, José M.<sup>a,b</sup>

<sup>a</sup> Departamento de Química Orgánica. Facultad de Ciencias, Universidad de Alicante, 03080-Alicante, Spain. Centro de Innovación en Química Avanzada (ORFEO-CINQA).

<sup>b</sup> Instituto de Síntesis Orgánica (ISO). Universidad de Alicante, 03080-Alicante, Spain.

\* cnajera@ua.es.



Dedicated to Prof. Dieter Enders


Received:  
Accepted:  
Published online:  
DOI:

**Abstract** The synthesis of polyfunctionalized indolizidines from pipecolinic acid alkyl ester derivatives, aldehydes and a wide range of dipolarophiles by a multicomponent 1,3-dipolar cycloadditions has been developed in a diastereoselective manner. Reactions take place in toluene with short reaction times at 70 °C, giving good yields. The synthesis of these fused heterocycles is also studied starting from the pipecolinic acid, generating the dipole through a decarboxylative route at 120 °C. The relative configuration of the resulting products, as well as the mechanistic pathways are also explained.

**Key words** cycloaddition, multicomponent, indolizidine, azomethine ylides, iminium route, decarboxylative route

## Introduction

The indolizidine structure can be found in many alkaloid families being the most important moiety in the molecule.<sup>1</sup> These alkaloids, which can be isolated from plant or animal sources, have shown important biological properties and medicinal applications.<sup>2</sup> As representative examples, pharmaceutically interesting tetrahydroxyalkaloids such as castanospermine and 6-epicastanospermine are possible lead compounds in the search for anti-AIDS drugs, and the most simple structure of tashiromine has multiple interesting biological activities (Figure 1).




**Figure 1.** Representative natural indolizidine alkaloids.

Many synthetic approaches have been successfully developed to prepare this fused heterocyclic skeleton. Most of them can be classified according to the cyclization order, that means, five-member ring followed by six-membered ring construction (5→6) and *vice versa* (6→5).<sup>3</sup> The most common used access is the 5→6 pathway, and the reason is the high availability of both natural or synthetic polyfunctionalized pyrrolidines or proline derivatives.<sup>4,5</sup> In addition, interesting 6→5 sequences have been published.<sup>6</sup>

Such as it has been extensively demonstrated, 1,3-dipolar cycloaddition (1,3-DC) involving azomethine ylides is a potential tool for the construction of complex alkaloid structures.<sup>7,8,9</sup> Focusing on 6→5 sequences mediated by these type of cycloadditions generating the intermediate azomethine ylides via decarboxylation route,<sup>10</sup> intramolecular 1,3-DC between  $\delta$ -chloroaldehydes, glycine and electrophilic alkenes, gave indolizidines in very high yields (Scheme 1a).<sup>11</sup> Another decarboxylations underwent by tetrahydroisoquinoline-3-carboxylic acid and by tetrahydro- $\beta$ -carboline-3-carboxylic acid also gave polycycles in good yields.<sup>12</sup>



The generation of the azomethine ylide through the iminium route, where a decarboxylation do not occur, has also been explored.<sup>13,14,15</sup> Ethyl pipecolate was used as starting material in the generation of the corresponding iminium salt only with benzaldehyde at 120 °C yielding mixture of stereoisomers when *N*-methylmaleimide, dimethyl fumarate and maleate were used (Scheme 1b).<sup>16</sup> In all these examples the chemical yield was almost quantitative (95%) and the mixture of diastereoisomers was notable. In the case of dimethyl fumarate and *N*-phenylmaleimide (NPM) a 50:50 and a 75:25 mixture of *endo:exo* adducts was obtained, respectively. The intrinsic thermal isomerization of dimethyl maleate also promoted the generation of a third diastereoisomer (Scheme 1b).



Following with the multicomponent 1,3-dipolar cycloaddition strategy designed for the synthesis of pyrrolizidines<sup>17</sup> using proline ester hydrochlorides, aldehydes and dipolarophiles, at room temperature, we will survey in this work the general scope of this cycloaddition employing six-membered ring templates for the construction of the fleeting azomethine ylide with aldehydes, and further capture with dipolarophiles. We will study the generation of these intermediates through the iminium route, or through the decarboxylation way. All these reactions will be designed with the idea of increasing the functionality of the resulting polysubstituted indolizidines improving the diastereomeric ratio at the end of the processes. A comparison between both methodologies will be also established.



## Results and discussion

Following the optimized reaction conditions described by our group in the synthesis of pyrrolizidines,<sup>17a,c</sup> the first attempt to run 1,3-DC between the iminium salt, generated from methyl pipercolinate hydrochloride **1a** and cinnamaldehyde, and further cyclization with *N*-substituted maleimides as dipolarophiles was performed at 25 °C in toluene. The reaction did not take place neither using these reaction conditions nor employing the silver catalyzed-process at room temperature. When increasing the temperature to 70 °C in toluene and in the absence of silver salts the reaction succeeded. Indolizidines **2** and **3** were isolated with high *endo*-diastereoselectivity when methyl pipercolinate and one equivalent of triethylamine were allowed to react with *N*-methylmaleimide (NMM) or with *N*-phenylmaleimide (NPM). No differences were observed concerning both of the yield the diastereoselectivity (70:30 *dr*). Compound **2** was isolated in 81% overall yield (*endo*-**2** 59%, and *exo*-**2** 22%) and **3** in 75% overall yield (*endo*-**3** 55%, and *exo*-**3** 20%) (Scheme 2). In contrast, when maleic anhydride was tested higher yield and complete *endo*-diastereoselectivity was achieved for compound **4** (Figure 2).




Following with the study of the general scope several dipolarophiles maintaining the aldehyde structure were tested. The highest *endo*-diastereoselection was achieved when the less sterically hindered methyl acrylate was employed as dipolarophile (Scheme 3) yielding *endo*-**5** (57%) in a 95:5 *dr*. When methyl fumarate was used as dipolarophile, the diastereoselectivity observed for compound *endo*-**6** (84:16 *dr*) was slightly lower than the obtained one for the methyl acrylate but higher than with NMM (Scheme 3). *trans*- $\beta$ -Nitrostyrene and chalcone were suitable dipolarophiles for this thermal multicomponent reaction affording indolizidines **7**, **8** and **9**, respectively. With the conjugated ketone the chemical yield was lower (66% overall yield) than the analogous obtained in the example run with the nitroalkene (72% overall yield). However, apart from the *endo*-**7** cycloadduct (17% yield), the regioisomer *exo*-**8** (55% yield) was also identified in high proportions. In the case of chalcone the unexpected cycloadduct *exo*-**9** was obtained as major product (75:25 *dr*) (Scheme 3).

Methyl 1,2,3,4-tetrahydroisoquinoline-3-carboxylate **10**, prepared from phenyl alanine methyl ester<sup>18</sup> was also tested in this multicomponent process. Cinnamaldehyde, **10** and NPM were mixed and the reaction was warmed at 70 °C for 17 h. Tetracyclic complex skeleton *endo*-**11** was obtained as major compound in 65% yield, and *exo*-**11** in 11% yield with a 86:14 *dr* (Scheme 4).



A comparison with the thermal 1,3-DC reported in the literature by Joucla and coworkers (Scheme 1b),<sup>16</sup> where benzaldehyde was employed, was done. The reaction with *N*-methylmaleimide did not work under our reaction conditions, and a complex mixture of inseparable products was detected after performing the reaction at 110 °C. However, the reaction involving maleic anhydride, benzaldehyde, and **1a** at 70 °C gave a modest isolated chemical yield (27%) of cycloadduct *endo*-**12** as unique diastereoisomer (Scheme 5). Furfural was also tested together with ethyl pipecolinate **1b** and NMM affording *endo*-cycloadduct **13** as major compound in 82:18 *dr* and 48% overall yield (Scheme 5).


Nevertheless, crotonaldehyde, isovaleraldehyde and (2*E*,4*E*)-hexa-2,4-dienal showed very poor reactivity. According to our studies, reactions run with all type of aldehydes, but specially unsaturated aldehydes, were very sensitive to high temperatures (>70 °C) affording decomposition products (detected in crude <sup>1</sup>H-NMR spectra).




After careful analysis of <sup>1</sup>H-NMR of the reaction crudes, only a mixture of two diastereoisomers was identified. At this point, we can assume that the reaction mechanism proceeds with relative high to excellent diastereocontrol as consequence of the S-type dipole generated by iminium route which attacks preferentially by its α-position to carbon atom of dipolarophile with partial positive charge (Figure 3a). This S-dipole, through the most favorable *endo*-approach allows the 2,4-*trans*-2,5-*trans* arrangement of the five-membered ring (Figure 3b). This

preferential trend was observed in compounds **2**, **3**, **4**, **5**, **6**, **11**, **12** and **13**. Only in the example run with the chalcone the *exo*-diastereoselectivity was preferentially observed presumably due to the high steric interaction between the phenyl group closer to ketone group and the substituents of the cyclic dipole.<sup>19</sup> In this example a 2,4-*cis*-2,5-*trans* arrangement was generated. On the other hand, the cycloadduct resulting from the  $\gamma$ -attack of the S-dipole was preferred in the reaction performed with *trans*- $\beta$ -nitrostyrene, which is an excellent Michael acceptor able to trap whatever type of resonance forms. In this case, the new compound **8** was formed by a more feasible *exo*-attack of the dipolarophile. This identical behavior was described during the multicomponent synthesis of pyrrolizidines because of stereoelectronic effects of the nitroalkene.<sup>17</sup>

The relative configuration of the major *endo*-cycloadducts **2-6** and **11-13** was proposed according to unambiguous nOe experiments. A representative example of these analyses is shown in Figure 4. The most stable conformations of *endo*-**5** and *endo*-**6** have been represented with the corresponding nOe effects detected. It is noteworthy to indicate that a small, but definitive, increment of electronic population in the signal of methyl group was produced when the hydrogen atom bonded to the C3 position was irradiated. In addition, this configuration was identical to the obtained by Joucla *et al.* after X-ray diffraction analysis,<sup>16</sup> and also with the same structural arrangement of pyrrolizidines obtained after multicomponent 1,3-DC involving prolinates.<sup>17</sup> At the end of the experimental work an X-ray diffraction analysis of crystalline major diastereoisomer *endo*-**13**<sup>20</sup> could be performed confirming all these stereochemical results obtained by NMR experiments (Scheme 5).



**Figure 3.** Mechanistic details of the synthesis of indolizidines by the iminium route.




**Figure 4.** Representative nOe detected for major *endo*-adducts (**4** and **5**) of this series of reactions.

Next, we study the availability of performing the corresponding 1,3-DC starting from pipercolinic acid **14**, aldehydes and dipolarophiles. In this reaction, the necessary decarboxylation of the iminium salt generated in situ occurred at higher temperatures (refluxing toluene) (Scheme 6). To the best of our knowledge, this type of cycloaddition of **14** has not been reported yet.<sup>21</sup> Thus, compound **14**, cinnamaldehyde, and NMM were diluted in toluene and the mixture was heated in a sealed tube at 120 °C (bath temperature) obtaining a mixture of four different stereoisomers **15** in 89% overall yield (Table 1, entry 1). In all cases the diastereomeric ratio observed in the crude <sup>1</sup>H-NMR spectra was identical to the analogous one determined after separation of each isomers by column chromatography. With NPM chemical yield of **16** was lower (78%) but the *endo*-diastereoselection was the highest of all this series of decarboxylative reactions using cinnamaldehyde (Table 1, entry 2). Dimethyl and diisobutyl fumarates gave both identical chemical yields (75%) and mixtures of diastereoisomers of products **17** and **18** (Table 1, entries 3 and 4). Non-symmetrical dipolarophiles such as *tert*-butyl acrylate and *trans*- $\beta$ -nitrostyrene were next evaluated. Compounds **19** and **20** were isolated in 52 and 40% overall yields, respectively (Table 1, entries 5 and 6). Finally, benzaldehyde was tested with **14** and NPM affording a very high yield (95%) of compound **21** as mixture of four stereoisomers (Table 1, entry 7). However, the reaction with NMM or *tert*-butyl acrylate only afforded decomposition products at the end of the reaction. Other aldehydes such as crotonaldehyde, isovaleraldehyde and furfural also failed as starting materials in the multicomponent reaction employing NMM.

After careful analyses of selective nOe experiments of each isolated product/mixture, and by comparison of the analogous chemical shifts and coupling constants we could identify each structure. *Endo*-cycloadduct was always the most abundant stereoisomer, followed by the *exo*-adduct and a couple of similar diastereoisomers with different olefinic chemical shifts. nOe experiments revealed clear *all-cis*-arrangement in C2, C3, C4, and C5, for the *endo*-cycloadduct **15** (Figure 5). Apart from *cis*-maleimide ring junction nOe, a very small one was observed between hydrogens bonded to C2 and C5 positions (Figure 5). A crucial nOe between hydrogens placed at C2 and the closer olefinic one was observed in both *endo*'- and *exo*'-**15** (Figure 5).

With all these stereochemical information, we can propose that the reaction, once produced the decarboxylation, operated through the most stable S-shape dipole interacting with the electrophilic alkene by its  $\alpha$ -position. S-Shape dipole underwent thermal stereomutation in the iminium salt affording unstable U-shape dipole, which is the responsible of the generation of

*endo'*- and *exo'*-diastereoisomers. Besides, the regioselectivity of this cycloaddition was very high because we could not detect any stereoisomer due to the attack of the dipole by its  $\gamma$ -position (Figure 6).




**Table 1.** Multicomponent 1,3-DC by mixing amino acid **14**, aldehydes, and dipolarophiles at 120 °C.

Entry	R <sup>1</sup>	Dipolarophile	Product	Yield (%) <sup>a</sup>	<i>dr</i> <sup>b</sup> ( <i>endo</i> : <i>exo</i> : <i>endo'</i> : <i>exo'</i> )
1	( <i>E</i> )-PhCH=CH-	NMM	<b>15</b>	89	35:22:20:23
2	( <i>E</i> )-PhCH=CH-	NPM	<b>16</b>	78	45:17:18:20
3	( <i>E</i> )-PhCH=CH-	Dimethyl fumarate	<b>17</b>	75	33:29:18:20
4	( <i>E</i> )-PhCH=CH-	Diisobutyl fumarate	<b>18</b>	75	35:30:19:17
5	( <i>E</i> )-PhCH=CH-	<i>tert</i> -Butyl acrylate	<b>19</b>	52	39:28:17:16
6	( <i>E</i> )-PhCH=CH-	<i>trans</i> - $\beta$ -Nitrostyrene	<b>20</b>	40	43:25:11:21
7	Ph	NPM	<b>21</b>	95	57:25:13:5


<sup>a</sup> Overall chemical yield isolated after flash chromatography.

<sup>b</sup> Mixture of diastereoisomers detected by <sup>1</sup>H-NMR of the crude mixture and after the separation of all of the corresponding diastereoisomers.



**Figure 5.** Representative nOe effects detected for adducts **15** derived from pipercolinic acid **14** and NMM.

As conclusion, we have prepared indolizidines from methyl pipercolinate and methyl 1,2,3,4-tetrahydroisoquinoline-3-carboxylate in a multicomponent 1,3-DC using functionalized conjugate aldehydes as cinnamaldehyde and different dipolarophiles at 70 °C. The major isomers resulted from the attack of the reactive S-shape dipole, prepared via the iminium route, by its  $\alpha$ -position affording *all cis-endo* diastereoisomers. Under these reaction conditions, the diastereoselection was notably higher than the analogous ones reported in the literature at 110 °C. In other side, the appearance of the U-shape dipole at 120 °C allowed to obtain two more diastereoisomers (*endo'* and *exo'*) when the multicomponent sequence dealt with a dipole generated by a decarboxylative route. Here, *endo*-cycloadduct was the major isomer, especially when a bulky substituent in the dipolarophile was bonded (Ph, Bu<sup>t</sup>, Bu<sup>i</sup>) but with very significant amounts of the corresponding *exo*-adduct. In all these examples, the diastereomeric ratio was very low. No regioisomeric adducts controlled by a  $\gamma$ -attack were obtained in any case.





## Experimental part

Melting points were determined with a Reichert Thermovar hot plate apparatus and are uncorrected. Only the structurally most important peaks of the IR spectra, recorded on a Nicolet 510 P-FT, are listed. For solid samples ATR device was employed. <sup>1</sup>H-NMR (300 MHz) and <sup>13</sup>C-NMR (75 MHz) spectra were obtained on a Bruker AC-300 using CDCl<sub>3</sub> as solvent and TMS as internal standard.

Low-resolution electron impact (EI) mass spectra were obtained at 70 eV on a Shimadzu QP-5000 and high-resolution mass spectra were obtained on a Finnigan VG Platform. HRMS (EI) were recorded on a Finnigan MAT 95S. Analytical TLC was performed on Schleicher & Schuell F1400/LS silicagel plates and the spots were visualized under UV light (λ = 254 nm). Flash chromatography was done using Merck silica gel 60 (0.040–0.063 mm).

## General procedure for the synthesis of indolizidines 3-9

To a solution of the pipercolic acid methyl ester hydrochloride **1** (40 mg, 0.22 mmol) in toluene (1 mL), Et<sub>3</sub>N (1 equiv, 30.5 μL, 0.22 mmol), the corresponding aldehyde (1 equiv, 0.22 mmol) and the dipolarophile (1 equiv, 0.22 mmol) were added. The resulting mixture was stirred at 70 °C for 17 h. EtOAc (5 mL) and H<sub>2</sub>O (5 mL) were added and the organic phase was separated, dried (MgSO<sub>4</sub>), and evaporated to obtain the crude heterocycle, which was purified by flash chromatography (silica-gel) in chemical yields reported along the text.

**Methyl (3aS\*,4S\*,9aR\*,9bR\*)-2-methyl-1,3-dioxo-4-[(E)-styryl]decahydro-9aH-pyrrolo[3,4-a]indolizine-9a-carboxylate (endo-2)**

Yield: 36 mg (59%), colorless prisms, mp 134–135 °C (Et<sub>2</sub>O).

IR (neat): 1734, 1698, 1213 cm<sup>-1</sup>.

<sup>1</sup>H-NMR: δ = 1.18 (dt, *J* = 13.3, 3.5 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.27–1.48 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.45–1.63 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>), 1.74 (dt, *J* = 13.2, 3.4 Hz, 1H, CCH<sub>2</sub>), 2.48 (ddd, *J* = 13.2, 2.9, 1.4 Hz, 1H, CCH<sub>2</sub>), 2.81, 2.84 (2xd, *J* = 2.7 Hz, 2H, NCH<sub>2</sub>), 3.01 (s, 3H, NCH<sub>3</sub>), 3.25 (dd, *J* = 8.0, 7.9 Hz, 1H, NCHCH), 3.35 (d, *J* = 7.9 Hz, 1H, CCH), 3.76 (s, 3H, OCH<sub>3</sub>), 4.18 (dd, *J* = 9.2, 8.0 Hz, 1H, NCH), 5.88 (dd, *J* = 15.6, 9.2 Hz, 1H, PhCHCH), 6.68 (d, *J* = 15.6 Hz, 1H, PhCH), 7.22–7.35 (m, 3H, ArH), 7.36–7.45 (m, 2H, ArH).

<sup>13</sup>C-NMR: δ = 21.7 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.7 (NCH<sub>2</sub>CH<sub>2</sub>), 25.1 (NCH<sub>3</sub>), 32.0 (CCH<sub>2</sub>), 43.8 (NCH<sub>2</sub>), 47.9 (NCHCHCO), 51.4 (CCHCO), 51.8 (OCH<sub>3</sub>), 65.2 (NCH), 69.9 (CCO<sub>2</sub>Me), 126.7, 127.8, 128.6, 128.7, 134.5, 136.8 (ArC, C=C), 173.6, 175.3 (2xNCO), 175.9 (CO<sub>2</sub>Me).

MS (EI): *m/z* = 368 (M<sup>+</sup>, 3%), 310 (20), 309 (100), 224 (3).

HRMS (DIP) calcd. for C<sub>21</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub>: 368.1746; found 368.1750.

**Methyl (3aS\*,4S\*,9aR\*,9bR\*)-1,3-dioxo-2-phenyl-4-[(E)-styryl]decahydro-9aH-pyrrolo[3,4-a]indolizine-9a-carboxylate (endo-3)**

Yield: 52 mg (55%), pale yellow oil.

IR (neat): 2933, 1710, 1498, 1448, 1375, 1179, 1309, 1192 cm<sup>-1</sup>.

<sup>1</sup>H-NMR: δ = 1.17–1.23 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.28–1.35 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.38–1.48 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.62–1.68 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.78 (dt, *J* = 13.1, 3.2 Hz, 1H, CCH<sub>2</sub>), 2.53 (dd, *J* = 13.1, 1.5 Hz, 1H, CCH<sub>2</sub>), 2.82–2.93 (m, 2H, NCH<sub>2</sub>), 3.41 (t, *J* = 8.0 Hz, 1H, NCHCH), 3.51 (d, *J* = 8.0 Hz, 1H, CCH), 3.79 (s, 3H, OCH<sub>3</sub>), 4.29 (dd, *J* = 8.9, 8.0 Hz, 1H, NCH), 6.01 (dd, *J* = 15.7, 8.9 Hz, 1H, PhCHCH), 6.72 (d, *J* = 15.7 Hz, 1H, PhCH), 7.16–7.35 (m, 5H, ArH), 7.37–7.50 (m, 5H, ArH).

<sup>13</sup>C-NMR: δ = 21.8 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.9 (NCH<sub>2</sub>CH<sub>2</sub>), 32.1 (CCH<sub>2</sub>), 44.0 (NCH<sub>2</sub>), 48.0 (NCHCHCO), 51.4 (CCHCO), 51.9 (OCH<sub>3</sub>), 65.5 (NCH), 70.4 (CCO<sub>2</sub>Me), 126.7, 126.9, 127.7, 127.9, 128.6, 128.8, 129.3, 132.0, 134.4, 136.8 (ArC, C=C), 173.6 (CO), 174.2 (CO), 175.0 (CO<sub>2</sub>Me).

MS (EI): *m/z* = 430 (M<sup>+</sup>, 3%), 372 (26), 371 (100), 224 (6).

HRMS (DIP) calcd. for C<sub>26</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub>: 430.1893; found 430.1911.

**Methyl (3aS\*,4S\*,9aR\*,9bR\*)-1,3-dioxo-4-[(E)-styryl]octahydro-3H,9aH-furo[3,4-a]indolizine-9a-carboxylate (endo-4)**

Yield: 72 mg (93%), colorless prisms, mp 146–148 °C (Et<sub>2</sub>O).

IR (neat): 1774, 1734, 1226 cm<sup>-1</sup>.

<sup>1</sup>H-NMR: δ = 1.15–1.25 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.50 (ddt, *J* = 12.9, 8.4, 3.9 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.58–1.69 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.67–1.85 (m, 2H, CCH<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>), 2.45 (ddd, *J* = 12.7, 4.5, 1.9 Hz, 1H, CCH<sub>2</sub>), 2.90 (d, *J* = 2.7 Hz, 1H, NCH<sub>2</sub>), 2.93 (d, *J* = 2.7 Hz, 1H, NCH<sub>2</sub>), 3.54 (dd, *J* = 8.5, 8.3 Hz, 1H, NCHCH), 3.71 (d, *J* = 8.5 Hz, 1H, CCH), 3.79 (s, 3H, OCH<sub>3</sub>), 4.27 (dd, *J* = 9.3, 8.3 Hz, 1H, NCH), 5.98 (dd, *J* = 15.7, 9.3 Hz, 1H, PhCHCH), 6.73 (d, *J* = 15.7 Hz, 1H, PhCH), 7.26–7.36 (m, 3H, ArH), 7.39–7.44 (m, 2H, ArH).

<sup>13</sup>C-NMR: δ = 21.3 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.3 (NCH<sub>2</sub>CH<sub>2</sub>), 31.5 (CCH<sub>2</sub>), 44.1 (NCH<sub>2</sub>), 48.4 (NCHCHCO), 52.0 (CCHCO), 52.4 (OCH<sub>3</sub>), 66.0 (NCH), 70.6 (CCO<sub>2</sub>Me), 125.5, 127.1, 128.4, 128.8, 136.2, 136.4 (ArC, C=C), 169.0, 169.4 (2xNCO), 172.4 (CO<sub>2</sub>Me).

MS (EI): *m/z* = 355 (M<sup>+</sup>, 5%), 297 (20), 296 (100), 225 (10), 224 (50).

HRMS (DIP) calcd. for C<sub>20</sub>H<sub>21</sub>NO<sub>5</sub>: 355.1420; found 355.1426.

**Dimethyl (2S\*,3S\*,8aR\*)-3-[(E)-styryl]hexahydroindolizine-2,8a(1H)-dicarboxylate (endo-5)**

Yield: 43 mg (57%), yellow solid, mp 189–190 °C (Et<sub>2</sub>O).

IR (neat): 2977, 2946, 1745, 1474 cm<sup>-1</sup>.

<sup>1</sup>H-NMR: δ = 1.11–1.17 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.33–1.51 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.52–1.60 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.65–1.75 (m, 1H, CCH<sub>2</sub>), 2.15 (dd, *J* = 12.4, 8.0 Hz, 1H, CHCH<sub>2</sub>), 2.24 (dd, *J* = 12.4, 10.8 Hz, 1H, CHCH<sub>2</sub>), 2.40 (dtd, *J* = 12.4, 3.3, 1.2 Hz, 1H, CCH<sub>2</sub>), 2.68 (td, *J* = 11.7, 3.3 Hz, 1H, NCH<sub>2</sub>), 2.80 (dd, *J* = 11.7, 3.9 Hz, 1H, NCH<sub>2</sub>), 3.20 (td, *J* = 10.5, 8.0 Hz, 1H, NCHCH), 3.53 (s, 3H, CHCO<sub>2</sub>CH<sub>3</sub>), 3.70 (s, 3H, CCO<sub>2</sub>CH<sub>3</sub>), 4.10 (ddd, *J* = 10.8, 8.8, 8.0 Hz, 1H, NCH), 5.91 (dd, *J* = 15.8, 8.8 Hz, 1H, PhCHCH), 6.53 (d, *J* = 15.8 Hz, 1H, PhCH), 7.15–7.21 (m, 1H, ArH), 7.23–7.30 (m, 2H, ArH), 7.31–7.38 (m, 2H, ArH).

<sup>13</sup>C-NMR: δ = 22.0 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 25.3 (NCH<sub>2</sub>CH<sub>2</sub>), 34.6 (CCH<sub>2</sub>CH<sub>2</sub>), 39.0 (CCH<sub>2</sub>CH), 43.4 (NCH<sub>2</sub>), 45.5 (CHCO<sub>2</sub>Me), 51.2 (OCH<sub>3</sub>), 51.6 (OCH<sub>3</sub>), 65.2 (NCH), 68.6 (CCO<sub>2</sub>Me), 126.4, 127.5, 128.5, 129.7, 132.9, 136.9 (ArC, C=C), 173.3 (CO<sub>2</sub>Me), 175.6 (CO<sub>2</sub>Me).

MS (EI): *m/z* = 343 (M<sup>+</sup>, 2%), 285 (20), 284 (100), 224 (12).

HRMS (DIP) calcd. for C<sub>20</sub>H<sub>25</sub>NO<sub>4</sub>: 343.1784; found: 343.1800.

**Trimethyl (1S\*,2S\*,3S\*,8aR\*)-3-[(E)-styryl]hexahydroindolizine-1,2,8a(1H)-tricarboxylate (endo-6)**

Yield: 62 mg (70%), pale yellow oil.

IR (neat): 1732, 1201, 1167 cm<sup>-1</sup>.

<sup>1</sup>H-NMR: δ = 1.24 (tdd, *J* = 13.5, 8.8, 3.9 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.46–1.65 (m, 3H, NCH<sub>2</sub>CH<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.68–1.79 (m, 1H, CCH<sub>2</sub>), 2.40–2.72 (m, 2H, NCH<sub>2</sub>, CCH<sub>2</sub>), 2.76–2.85 (m, 1H, NCH<sub>2</sub>), 3.43 (d, *J* = 10.8 Hz, 1H, CCH), 3.55 (s, 3H, CHCO<sub>2</sub>CH<sub>3</sub>), 3.68 (s, 3H, CHCO<sub>2</sub>CH<sub>3</sub>), 3.69 (dd, *J* = 10.8, 10.5 Hz, 1H, NCHCH), 3.70 (s, 3H, CCO<sub>2</sub>CH<sub>3</sub>), 4.19 (dd, *J* = 10.5, 9.1 Hz, 1H, NCH), 5.87 (dd, *J* = 15.8, 9.1 Hz, 1H, PhCHCH), 6.55 (d, *J* = 15.8 Hz, 1H, PhCH), 7.19–7.37 (m, 5H, ArH).

<sup>13</sup>C-NMR: δ = 21.8 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 25.1 (NCH<sub>2</sub>CH<sub>2</sub>), 34.5 (CCH<sub>2</sub>), 43.8 (NCH<sub>2</sub>), 48.2 (CCH), 51.7 (NCHCH), 52.1 (OCH<sub>3</sub>), 52.3 (OCH<sub>3</sub>), 55.0 (OCH<sub>3</sub>), 64.4 (NCH), 70.9 (CCO<sub>2</sub>Me), 126.6, 127.8, 128.7, 128.9, 133.7, 136.8 (ArC, C=C), 171.0 (CCO<sub>2</sub>Me), 172.3 (CHCO<sub>2</sub>Me), 172.8 (CHCO<sub>2</sub>Me).

MS (EI): *m/z* = 401 (M<sup>+</sup>, 5%), 343 (21), 342 (100), 310 (13), 282 (38), 250 (23).

HRMS (DIP) calcd. for C<sub>22</sub>H<sub>27</sub>NO<sub>6</sub>: 401.1838; found 401.1849.

**Methyl (1S\*,2S\*,3S\*,8aR\*)-2-nitro-1-phenyl-3-[(E)-styryl]hexahydroindolizine-8a(1H)-carboxylate (endo-7)**

Yield: 15 mg (17%), brown sticky oil.

IR (neat): 1734, 1556, 1223 cm<sup>-1</sup>.

<sup>1</sup>H-NMR: δ = 1.21–1.38 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.50–1.70 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>), 1.74–1.87 (m, 1H, CCH<sub>2</sub>), 2.24–2.31 (m, 1H, CCH<sub>2</sub>), 2.96–3.02 (m, 2H, NCH<sub>2</sub>), 3.86 (s, 3H, OCH<sub>3</sub>), 3.93 (dd, *J* = 8.4, 5.2 Hz, 1H, NCHCHPh), 4.08 (t, *J* = 8.4 Hz, 1H, NCH), 5.22 (d, *J* = 5.2, 1H, CHNO<sub>2</sub>), 6.15 (dd, *J* = 15.8, 8.4 Hz, 1H, PhCHCH), 6.40 (d, *J* = 15.8 Hz, 1H, PhCH), 7.06–7.12 (m, 2H, ArH), 7.22–7.39 (m, 8H, ArH).



<sup>13</sup>C-NMR:  $\delta$  = 21.6 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.6 (NCH<sub>2</sub>CH<sub>2</sub>), 31.5 (CCH<sub>2</sub>), 43.7 (NCHCHPh), 52.4 (NCH<sub>2</sub>), 54.5 (OCH<sub>3</sub>), 71.5 (NCH), 72.0 (CCO<sub>2</sub>Me), 97.9 (CHNO<sub>2</sub>), 126.7, 127.7, 128.0, 128.2, 128.7, 129.1, 129.5, 134.7, 136.5, 138.1 (ArC, C=C), 172.8 (CO<sub>2</sub>Me).

MS (EI):  $m/z$  = 360 (M<sup>+</sup>-NO<sub>2</sub>, 91%), 348 (13), 347 (51), 302 (25), 301 (100), 300 (39), 224 (18), 210 (45), 198 (13).

HRMS (DIP) calcd. for C<sub>24</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub>(-NO<sub>2</sub>): 360.1983; found: 360.1974.

**Methyl (1S\*,2R\*,3S\*,8aR\*)-1-nitro-2-phenyl-3-[(E)-styryl]hexahydroindolizidine-8a(1H)-carboxylate (exo-8)**

Yield: 62 mg (55%), colorless prisms, mp 146–148 °C (Et<sub>2</sub>O).

IR (neat): 1543, 1354 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.08–1.32 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.47–1.65 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>), 1.69–1.82 (m, 1H, CCH<sub>2</sub>), 2.21–2.56 (m, 2H, NCH<sub>2</sub>, CCH<sub>2</sub>), 2.86–2.93 (m, 1H, NCH<sub>2</sub>), 3.43 (s, 3H, OCH<sub>3</sub>), 4.09 (d,  $J$  = 10.0 Hz, 1H, CCHPh), 4.69 (dd,  $J$  = 9.6, 8.4 Hz, 1H, NCH), 5.74 (dd,  $J$  = 10.0, 9.6 Hz, 1H, CHNO<sub>2</sub>), 5.90 (dd,  $J$  = 15.8, 8.4 Hz, 1H, PhCHCH), 6.72 (d,  $J$  = 15.8 Hz, 1H, PhCH), 7.08–7.21 (m, 2H, ArH), 7.27–7.34 (m, 6H, ArH), 7.35–7.42 (m, 2H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 21.9 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 25.1 (NCH<sub>2</sub>CH<sub>2</sub>), 33.6 (CCH<sub>2</sub>), 44.6 (CCHPh), 51.1 (NCH<sub>2</sub>), 57.9 (OCH<sub>3</sub>), 65.7 (NCH), 73.4 (CCO<sub>2</sub>Me), 91.1 (CHNO<sub>2</sub>), 126.0, 127.0, 127.9, 127.9, 128.2, 128.4, 128.6, 128.8, 133.5, 136.3 (ArC, C=C), 173.6 (CO<sub>2</sub>Me).

MS (EI):  $m/z$  = 360 (M<sup>+</sup>-NO<sub>2</sub>, 57%), 348 (19), 347 (80), 301 (33), 300 (100).

HRMS (DIP) calcd. for C<sub>24</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub>(-NO<sub>2</sub>): 360.1983; found: 360.1974.

**Methyl (1R\*,2S\*,3S\*,8aR\*)-2-benzoyl-1-phenyl-3-[(E)-styryl]hexahydroindolizidine-8a(1H)-carboxylate (endo-9) and Methyl (1S\*,2R\*,3S\*,8aR\*)-2-benzoyl-1-phenyl-3-[(E)-styryl]hexahydroindolizidine-8a(1H)-carboxylate (exo-9)**

Yield: 68 mg (66%), colorless sticky oil.

IR (neat): 1718, 1682, 1447, 1207 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  (mixture of *endo:exo* 0.33:1) = 1.13–1.19 (m, *exo*-2H, NCHCHCH<sub>2</sub>), 1.20–1.24 (m, *endo*-2H, NCHCHCH<sub>2</sub>), 1.49–1.56 (m, *endo*-2H, NCHCH<sub>2</sub>, *exo*-2H, NCHCH<sub>2</sub>), 1.68–1.81 (m, *endo*-1H, CCH<sub>2</sub>, *exo*-1H, CCH<sub>2</sub>), 1.88–1.93 (m, *endo*-2H, NCH<sub>2</sub>, CCH<sub>2</sub>), 2.36 (dt,  $J$  = 12.3, 3.4 Hz, *exo*-1H, CCH<sub>2</sub>), 2.49 (td,  $J$  = 11.4, 4.0 Hz, *exo*-1H, NCH<sub>2</sub>), 2.87 (dd,  $J$  = 15.3, 3.8 Hz, *exo*-1H, NCH<sub>2</sub>), 2.96–2.92 (m, *endo*-1H, NCH<sub>2</sub>), 3.42 (s, *exo*-3H, OCH<sub>3</sub>), 3.90 (dd,  $J$  = 8.5, 6.3 Hz, *endo*-1H, NCHCH), 3.93 (s, *endo*-3H, OCH<sub>3</sub>), 4.03 (d,  $J$  = 11.2 Hz, *exo*-1H, CCH), 4.13 (dd,  $J$  = 8.5, 8.3 Hz, *endo*-1H, NCH), 4.26 (d,  $J$  = 6.3 Hz, *endo*-1H, CCH), 4.58 (dd,  $J$  = 10.2, 8.8 Hz, *exo*-1H, NCH), 4.85 (dd,  $J$  = 11.2, 10.2 Hz, *exo*-1H, NCHCH), 5.71 (dd,  $J$  = 15.7, 8.8 Hz, *exo*-1H, PhCHCH), 6.20 (dd,  $J$  = 15.9, 8.3 Hz, *endo*-1H, PhCHCH), 6.30 (d,  $J$  = 15.7 Hz, *exo*-1H, PhCH), 6.37 (d,  $J$  = 15.9 Hz, *endo*-1H, PhCH), 7.04–7.23 (m, *endo*-9H, *exo*-9H, ArH), 7.28–7.46 (m, *endo*-4H, *exo*-4H, ArH), 7.83–7.92 (m, *exo*-2H, ArH), 8.01–8.08 (m, *endo*-2H, ArH).

<sup>13</sup>C-NMR:  $\delta$  (major diastereoisomer) = 22.3 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 25.4 (NCH<sub>2</sub>CH<sub>2</sub>), 33.8 (CCH<sub>2</sub>), 44.0 (NCH<sub>2</sub>), 50.8 (CCHPh), 51.9 (CHCO), 55.2 (OCH<sub>3</sub>), 65.8 (NCH), 73.6 (CCO<sub>2</sub>Me), 126.5, 126.6, 127.4, 127.5, 128.1, 128.1, 128.3, 128.4, 128.5, 128.5, 128.6, 128.7, 128.9, 129.9, 133.0, 133.0, 136.7, 137.0, 138.1 (ArC, C=C), 174.6 (CO<sub>2</sub>Me), 198.5 (CO).

MS (EI):  $m/z$  = 465 (M<sup>+</sup>, 3%), 407 (30), 406 (100), 360 (8).

HRMS (DIP) calcd. for C<sub>31</sub>H<sub>31</sub>NO<sub>5</sub>: 465.2324; found 465.2334.

**General procedure for the synthesis of indolizidine 11**

To a solution of the amine **10** (40 mg, 0.21 mmol) in toluene (1 mL), the corresponding aldehyde (1 equiv, 0.21 mmol) and the dipolarophile (1 equiv, 0.21 mmol) were added. The resulting mixture was stirred at 70 °C for 17 h. The solvent was evaporated and the heterocycles were separated by flash chromatography (silica-gel) in good chemical yields (see text).

**Methyl (3aS\*,4S\*,11aR\*,11bR\*)-1,3-dioxo-2-phenyl-4-[(E)-styryl]-1,2,3,3a,4,6,11,11b-octahydro-11aH-pyrrolo[3',4':3,4]pyrrolo[1,2-b]isoquinoline-11a-carboxylate (endo-11)**

Yield: 68 mg (65%), white solid, mp 209–212 °C (Et<sub>2</sub>O).

IR (neat): 1703, 1494, 1396, 1203 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 2.98 (d,  $J$  = 16.8 Hz, 1H, CCH<sub>2</sub>), 3.44 (d,  $J$  = 8.0 Hz, 1H, CCHCO), 3.52 (d,  $J$  = 16.8 Hz, 1H, CCH<sub>2</sub>), 3.55 (dd,  $J$  = 8.2, 8.0 Hz, 1H, NCHCH), 3.72 (s, 3H, OCH<sub>3</sub>), 3.88 (dd,  $J$  = 8.6, 8.2 Hz, 1H, NCH), 3.95 (d,  $J$  = 18.1 Hz, 1H, NCH<sub>2</sub>), 4.31 (d,  $J$  = 18.1 Hz, 1H, NCH<sub>2</sub>), 6.22 (dd,  $J$  = 15.7, 8.6 Hz, 1H, PhCHCH), 6.62 (d,  $J$  = 15.7 Hz, 1H, PhCH), 6.94–7.06 (m, 1H, ArH), 7.09–7.21 (m, 3H, ArH), 7.23–7.32 (m, 5H, ArH), 7.37–7.48 (m, 5H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 30.2 (CCH<sub>2</sub>), 45.4 (NCH<sub>2</sub>), 47.4 (NCHCHCO), 52.8 (CCHCO), 53.7 (OCH<sub>3</sub>), 64.8 (NCH), 68.8 (CCO<sub>2</sub>Me), 126.2, 126.5, 126.6, 126.9, 127.1, 128.2, 128.7, 128.8, 129.0, 129.2, 130.6, 131.9, 135.2, 136.3 (ArC, C=C), 170.9 (CO), 174.2 (CO), 174.6 (CO<sub>2</sub>Me).

MS (EI):  $m/z$  = 478 (M<sup>+</sup>, <1%), 420 (31), 419 (100), 180 (4).

HRMS (DIP) calcd. for C<sub>30</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub>: 478.1893; found 478.1883.

**General procedure for the synthesis of indolizidines 12–13**

To a solution of the pipecolic acid alkyl ester hydrochloride **1** (0.22 mmol) in toluene (1 mL), Et<sub>3</sub>N (1 equiv, 30.5  $\mu$ L, 0.22 mmol), the corresponding aldehyde (1 equiv, 0.22 mmol) and the dipolarophile (1 equiv, 0.22 mmol) were added. The resulting mixture was stirred at 70 °C for 17 h. EtOAc (5 mL) and H<sub>2</sub>O (5 mL) were added and the organic phase was separated, dried (MgSO<sub>4</sub>), and evaporated to obtain the crude product which was purified by flash chromatography (silica-gel) in good chemical yields (see text).

**Methyl (3aS\*,4R\*,9aR\*,9bR\*)-1,3-dioxo-4-phenyloctahydro-3H,9aH-furo[3,4-a]indolizidine-9a-carboxylate (endo-12)**

Yield: 20 mg (27%), yellow sticky oil.

IR (neat): 2927, 2856, 1781, 1733, 1209, 922, 734 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.53–1.39 (m, 2H, NCHCHCH<sub>2</sub>), 1.55–1.65 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.83 (dt,  $J$  = 13.8, 3.5 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.96 (td,  $J$  = 13.4, 3.9 Hz, 1H, CCH<sub>2</sub>), 2.45–2.55 (m, 1H, CCH<sub>2</sub>), 2.65 (dd,  $J$  = 11.9, 4.3 Hz, 1H, NCH<sub>2</sub>), 2.78 (td,  $J$  = 11.9, 3.3 Hz, 1H, NCH<sub>2</sub>), 3.61 (dd,  $J$  = 9.4, 8.3 Hz, 1H, PhCHCH), 3.69 (d,  $J$  = 8.3 Hz, 1H, CCHCO), 3.79 (s, 3H, OCH<sub>3</sub>), 4.75 (d,  $J$  = 9.4 Hz, 1H, PhCH), 7.22–7.28 (m, 2H, ArH), 7.29–7.40 (m, 3H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 21.4 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.6 (NCH<sub>2</sub>CH<sub>2</sub>), 31.0 (CCH<sub>2</sub>), 44.1 (NCH<sub>2</sub>), 49.6 (NCHCHCO), 52.2 (OCH<sub>3</sub>), 52.1 (CCHCO), 67.7 (NCH), 70.9 (CCO<sub>2</sub>Me), 127.8, 128.6, 128.8, 128.9, 130.2, 136.5 (ArC, C=C), 169.0, 169.2 (2xNCO), 172.9 (CO<sub>2</sub>Me).

MS (EI):  $m/z$  = 329 (M<sup>+</sup>, <1%), 271 (18), 270 (100), 220 (8), 198 (67).

HRMS (DIP) calcd. for C<sub>18</sub>H<sub>19</sub>NO<sub>5</sub>(-CO<sub>2</sub>Me): 270.1130; found 270.1132.

**Ethyl (3aS\*,4R\*,9aR\*,9bR\*)-4-(furan-2-yl)-2-methyl-1,3-dioxodecahydro-9aH-pyrrolo[3,4-a]indolizidine-9a-carboxylate (endo-13)**

Yield: 30 mg (39%), white prisms, mp 121–124 °C (Et<sub>2</sub>O).

IR (neat): 2936, 1699, 1432, 1377, 1281, 1230, 1148, 1006, 755 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.21 (dt,  $J$  = 13.3, 3.8 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.32 (t,  $J$  = 7.2 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>), 1.44 (ddd,  $J$  = 12.4, 5.3, 4.1 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.51–1.61 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.78 (dd,  $J$  = 13.9, 6.0 Hz, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.87 (td,  $J$  = 13.2, 3.7 Hz, 1H, CCH<sub>2</sub>), 2.53 (ddt,  $J$  = 13.2, 4.6, 2.1 Hz, 1H, CCH<sub>2</sub>), 2.56–2.65 (m, 1H, NCH<sub>2</sub>), 2.79 (td,  $J$  = 11.7, 3.5 Hz, 1H, NCH<sub>2</sub>), 2.93 (s, 3H, NCH<sub>3</sub>), 3.28–3.40 (m, 2H, NCHCH, CCH), 4.23 (q,  $J$  = 7.2 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 4.80 (d,  $J$  = 8.2 Hz, 1H, NCH), 6.24 (dd,  $J$  = 3.2, 0.8 Hz, 1H, OCHCHCH), 6.33 (dd,  $J$  = 3.2, 1.9 Hz, 1H, OCHCH), 7.38 (dd,  $J$  = 1.9, 0.8 Hz, 1H, OCH).

<sup>13</sup>C-NMR:  $\delta$  = 14.5 (CH<sub>2</sub>CH<sub>3</sub>), 21.6 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.7 (NCH<sub>2</sub>CH<sub>2</sub>), 25.1 (NCH<sub>3</sub>), 30.9 (CCH<sub>2</sub>), 44.3 (NCH<sub>2</sub>), 47.0 (NCHCHCO), 51.3 (CCHCO), 61.0 (CH<sub>2</sub>CH<sub>3</sub>), 61.2 (NCH), 69.9 (CCO<sub>2</sub>Et), 109.2 (OCHCH), 110.3 (OCCH), 142.8 (OCH), 151.1 (OCCH), 173.1, 175.0 (2xNCO), 175.8 (CO<sub>2</sub>Et).

MS (EI):  $m/z$  = 346 (M<sup>+</sup>, <1%), 274 (16), 273 (100).

HRMS (DIP) calcd. for C<sub>18</sub>H<sub>22</sub>N<sub>2</sub>O<sub>5</sub>: 346.1529; found: 346.1519.

**General procedure for the synthesis of indolizidines 15–21**

To a solution of the pipecolic acid **13** (40 mg, 0.31 mmol) in toluene (1 mL), the corresponding aldehyde (1 equiv, 0.31 mmol) and the dipolarophile (1 equiv, 0.31 mmol) were added. The resulting mixture was stirred in a pressure tube at 120 °C for 17 h. The solvent was evaporated and the mixture was separated by flash chromatography affording the corresponding cycloadducts.

**(3aS\*,4S\*,9aR\*,9bR\*)-2-Methyl-4-[(E)-styryl]octahydro-1H-pyrrolo[3,4-a]indolizine-1,3(2H)-dione (endo-15) and (3aR\*,4R\*,9aR\*,9bS\*)-2-Methyl-4-[(E)-styryl]octahydro-1H-pyrrolo[3,4-a]indolizine-1,3(2H)-dione (exo-15)**

Yield: 57 mg (59%), brown sticky oil.

IR (neat): 2938, 1697, 1433, 1382, 1281, 1239, 1138, 1039, 965, 749, 694 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  (mixture of *endo:exo'* 1:0.9) = 1.11–1.28 (m, *endo*-2H, NCHCH<sub>2</sub>, *exo'*-1H, NCHCH<sub>2</sub>), 1.34–1.45 (m, *endo*-1H, NCH<sub>2</sub>CH<sub>2</sub>, *exo'*-1H, NCHCH<sub>2</sub>), 1.50–1.66 (m, *endo*-1H, NCH<sub>2</sub>CH<sub>2</sub>, *exo'*-1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.69–1.90 (m, *endo*-1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, *exo'*-2H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>), 1.97–2.12 (m, *endo*-1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, *exo'*-1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.18–2.36 (m, *J* = m, *endo*-1H, NCH<sub>2</sub>, *exo'*-1H, NCH<sub>2</sub>), 2.79–2.93 (m, *endo*-2H, NCHCH<sub>2</sub>, NCH<sub>2</sub>, *exo'*-1H, NCHCH<sub>2</sub>), 2.98–3.02 (m, *endo*-3H, NCH<sub>3</sub>, *exo'*-4H, NCH<sub>3</sub>, NCHCH), 3.03–3.08 (m, *endo*-1H, NCHCH, *exo'*-1H, NCH<sub>2</sub>), 3.10–3.32 (m, *endo*-1H, CH<sub>2</sub>CHCH, *exo'*-2H, NCHCH, CH<sub>2</sub>CHCH), 4.13 (d, *J* = 9.6 Hz, *endo*-1H, NCHCH), 6.12 (dd, *J* = 15.7, 9.2 Hz, *exo'*-1H, PhCHCH), 6.26 (dd, *J* = 15.7, 9.6 Hz, *endo*-1H, PhCHCH), 6.61 (d, *J* = 15.7 Hz, *exo'*-1H, PhCH), 6.64 (d, *J* = 15.7 Hz, *endo*-1H, PhCH), 7.46–7.21 (m, *endo*-5H, *exo'*-5H, ArH).

<sup>13</sup>C-NMR:  $\delta$  (mixture of *endo:exo'*) = 24.4, 24.4 (2xNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.8, 25.0 (2xNCH<sub>2</sub>CH<sub>2</sub>), 25.0, 25.1 (2xNCH<sub>3</sub>), 28.0, 28.9 (2xNCHCH<sub>2</sub>), 47.1, 47.6, 48.2 (3xCHCO), 48.5 (NCH<sub>2</sub>), 50.0 (CHCO), 51.8 (NCH<sub>2</sub>), 60.5, 65.9, 67.8, 70.5 (4xNCH), 125.0, 126.5, 126.8, 127.9, 128.0, 128.2, 128.6, 128.7, 134.3, 134.5, 136.3, 136.7 (ArC, 2xC=C), 176.4, 176.7, 176.9, 178.7 (4xCO).

MS (EI): *m/z* = 310 (M<sup>+</sup>, 18%), 309 (17), 220 (14), 219 (100), 199 (20), 198 (17), 115 (10).

HRMS (DIP) calcd. for C<sub>19</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub>: 310.1681; found: 310.1668.

**(3aR\*,4S\*,9aR\*,9bS\*)-2-Methyl-4-[(E)-styryl]octahydro-1H-pyrrolo[3,4-a]indolizine-1,3(2H)-dione (exo-15)**

Yield: 21 mg (22%), yellow sticky oil.

IR (neat): 2919, 2850, 1698, 1435, 1283, 1122, 1074, 1010, 966, 732, 694 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.21–1.30 (m, 1H, NCHCH<sub>2</sub>), 1.44–1.53 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>, NCHCH<sub>2</sub>), 1.55–1.69 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.78–1.89 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.62–2.75 (m, 1H, NCH<sub>2</sub>), 2.88 (dd, *J* = 8.2, 2.1 Hz, 1H, CH<sub>2</sub>CHCH), 2.91–2.98 (m, 1H, NCH<sub>2</sub>), 3.00 (s, 1H, NCH<sub>3</sub>), 3.38 (dd, *J* = 8.2, 8.0 Hz, 1H, NCHCHCO), 3.44–3.51 (m, 1H, NCH<sub>2</sub>), 4.10 (dd, *J* = 9.5, 8.0 Hz, 1H, NCHCHCO), 5.93 (dd, *J* = 15.7, 9.5 Hz, 1H, PhCHCH), 6.64 (d, *J* = 15.7 Hz, 1H, PhCH), 7.23–7.42 (m, 5H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 19.2 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.5 (NCH<sub>2</sub>CH<sub>2</sub>), 25.2 (NCH<sub>3</sub>), 27.0 (NCHCH<sub>2</sub>), 45.5 (NCH<sub>2</sub>), 48.7 (NCHCHCO), 50.5 (NCHCHCO), 62.1, 62.1 (2xNCH), 126.5, 126.8, 127.9, 128.6, 134.6, 136.6 (ArC, C=C), 176.5, 178.8 (2xNCO).

MS (EI): *m/z* = 310 (M<sup>+</sup>, 18%), 309 (16), 220 (14), 219 (100), 199 (9), 198 (12), 115 (9).

HRMS (DIP) calcd. for C<sub>19</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub>: 310.1681; found: 310.1668.

**(3aS\*,4S\*,9aR\*,9bR\*)-2-Phenyl-4-[(E)-styryl]octahydro-1H-pyrrolo[3,4-a]indolizine-1,3(2H)-dione (endo-16)**

Yield: 31 mg (27%), yellow sticky oil.

IR (neat): 2941, 1708, 1498, 1381, 1185, 968, 849, 734 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.15–1.29 (m, 2H, NCHCH<sub>2</sub>CH<sub>2</sub>), 1.38–1.47 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.54–1.63 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>), 1.77–1.85 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.02–2.13 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.34 (td, *J* = 11.5, 3.0 Hz, 1H, NCH<sub>2</sub>), 2.85–2.92 (m, 1H, NCH<sub>2</sub>), 2.96 (ddd, *J* = 10.8, 8.4, 2.7 Hz, 1H, NCHCH<sub>2</sub>), 3.22 (dd, *J* = 7.9, 0.8 Hz, 1H, NCHCHCO), 3.43 (dd, *J* = 8.4, 7.9 Hz, 1H, CH<sub>2</sub>CHCH), 4.25 (d, *J* = 9.5 Hz, 1H, NCHCHCO), 6.30 (dd, *J* = 15.7, 9.5 Hz, 1H, PhCHCH), 6.67 (d, *J* = 15.7 Hz, 1H, PhCH), 7.19–7.52 (m, 10H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 24.5 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 25.2 (NCH<sub>2</sub>CH<sub>2</sub>), 29.3 (NCHCH<sub>2</sub>), 47.6 (NCHCHCO), 48.6 (NCH<sub>2</sub>), 50.2 (NCHCHCO), 60.8 (NCH), 68.2 (NCH), 124.8, 126.2, 126.9, 128.1, 128.6, 128.8, 129.1, 132.3, 134.4, 136.4 (ArC, C=C), 176.0, 177.8 (2xNCO).

MS (EI): *m/z* = 372 (M<sup>+</sup>, 23%), 371 (16), 282 (19), 281 (100), 199 (32), 198 (18), 115 (10).

HRMS (DIP) calcd. for C<sub>24</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>: 372.1838; found: 372.1828.

**(3aR\*,4S\*,9aR\*,9bS\*)-2-Phenyl-4-[(E)-styryl]octahydro-1H-pyrrolo[3,4-a]indolizine-1,3(2H)-dione (exo-16)**

Yield: 17 mg (15%), prisms, mp 133–137 °C (Et<sub>2</sub>O).

IR (neat): 2930, 1705, 1498, 1384, 1189, 974, 749 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.25–1.34 (m, 2H, NCHCH<sub>2</sub>), 1.46–1.62 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>), 1.69–1.76 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.79–1.95 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.66–2.79 (m, 1H, NCH<sub>2</sub>), 2.93–3.02 (m, 1H, NCHCH<sub>2</sub>), 3.06 (dd, *J* = 8.4, 2.4 Hz, 1H, CH<sub>2</sub>CHCH), 3.56 (dd, *J* = 8.4, 8.2 Hz, 1H, NCHCHCO), 3.53–3.59 (m, 1H, NCH<sub>2</sub>), 4.21 (dd, *J* = 9.1, 8.2 Hz, 1H, NCHCHCO), 6.05 (dd, *J* = 15.7, 9.1 Hz, 1H, PhCHCH), 6.69 (d, *J* = 15.7 Hz, 1H, PhCH), 7.23–7.46 (m, 10H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 19.7 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.5 (NCH<sub>2</sub>CH<sub>2</sub>), 27.4 (NCHCH<sub>2</sub>), 45.8 (NCH<sub>2</sub>), 48.7 (NCHCHCO), 50.6 (NCHCHCO), 62.5 (NCH), 62.8 (NCH), 125.9, 126.6, 126.9, 128.0, 128.6, 128.7, 129.2, 132.1, 134.7, 136.6 (ArC, C=C), 175.5, 177.8 (2xNCO).

MS (EI): *m/z* = 372 (M<sup>+</sup>, 23%), 371 (13), 282 (19), 281 (100), 199 (15), 198 (14).

HRMS (DIP) calcd. for C<sub>24</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>: 372.1838; found: 372.1828.

**Dimethyl (1S\*,2S\*,3S\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (endo-17) and dimethyl (1R\*,2R\*,3R\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (exo-17)**

Yield: 37 mg (35%), yellow oil.

IR (neat): 2934, 2853, 1733, 1436, 1300, 1196, 1168, 1011, 968, 749, 693 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  (mixture of *endo:exo'* 1:0.75, difficult assignment) = 1.04–1.41 (m, 4H), 1.49 (tt, *J* = 7.1, 3.6 Hz, 3H), 1.64–1.71 (m, 1H), 1.78 (td, *J* = 9.2, 7.4, 4.2 Hz, 2H), 1.87–1.96 (m, 1H), 2.36–2.50 (m, 2H), 2.80–2.96 (m, 3H), 3.08 (dd, *J* = 7.8, 7.1 Hz, 1H), 3.15 (ddd, *J* = 11.5, 8.9, 2.9 Hz, 1H), 3.27 (dd, *J* = 7.1, 4.2 Hz, 1H), 3.55 (s, 2H), 3.71 (s, *J* = 1.0 Hz, 5H), 3.75 (s, 2H), 3.88 (t, *J* = 8.1 Hz, 1H), 4.03–4.15 (m, 2H), 6.09 (dd, *J* = 15.7, 9.8 Hz, 1H), 6.28 (dd, *J* = 15.7, 9.5 Hz, 1H), 6.54 (d, *J* = 15.7 Hz, 1H), 6.57 (d, *J* = 15.7 Hz, 1H), 7.22–7.35 (m, 7H), 7.36–7.42 (m, 2H).

<sup>13</sup>C-NMR:  $\delta$  (mixture of *endo:exo'*, difficult assignment) = 23.4, 23.7, 24.0, 24.3, 27.3, 30.5, 47.6, 48.0, 48.1, 49.4, 51.5, 51.9, 52.0, 52.3, 52.3, 61.3, 63.3, 66.3, 67.2, 124.8, 126.6, 127.8, 127.8, 128.4, 132.2, 134.7, 136.6, 172.2, 173.1, 173.7, 173.8.

MS (EI): *m/z* = 343 (M<sup>+</sup>, 33%), 284 (35), 282 (19), 253 (15), 252 (100), 250 (17), 199 (39), 198 (24), 115 (17).

HRMS (DIP) calcd. for C<sub>20</sub>H<sub>25</sub>NO<sub>4</sub>: 343.1784; found: 343.1785.

**Dimethyl (1S\*,2S\*,3R\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (endo-17) and Dimethyl (1R\*,2R\*,3S\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (exo-17)**

Yield: 43 mg (40%), yellow oil.

IR (neat): 2944, 2854, 1733, 1436, 1196, 1167, 1005, 969, 746, 693 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  (mixture of *endo:exo'* 0.65:1, difficult assignment) = 1.06–1.25 (m, 2H), 1.43 (tdd, *J* = 12.4, 10.8, 3.6 Hz, 2H), 1.58 (q, *J* = 3.4 Hz, 2H), 1.76–1.91 (m, 4H), 1.98–2.07 (m, 1H), 2.19 (td, *J* = 10.4, 2.5 Hz, 1H), 2.43 (ddd, *J* = 10.8, 8.3, 2.3 Hz, 1H), 2.95–3.05 (m, 1H), 3.12 (d, *J* = 10.9 Hz, 1H), 3.18–3.32 (m, 3H), 3.37 (dd, *J* = 8.5, 4.9 Hz, 1H), 3.43 (dd, *J* = 10.0, 7.8 Hz, 1H), 3.59 (s, 3H), 3.66 (s, 2H), 3.70 (s, 3H), 3.73 (s, 2H), 5.95 (dd, *J* = 15.8, 8.6 Hz, 1H), 6.21 (dd, *J* = 15.8, 8.5 Hz, 1H), 6.55 (dd, *J* = 15.8 Hz, 1H), 6.57 (dd, *J* = 15.8 Hz, 1H), 7.16–7.45 (m, 9H).

<sup>13</sup>C-NMR:  $\delta$  (mixture of *endo:exo'*, difficult assignment) = 24.0, 24.3, 24.7, 24.9, 28.4, 30.5, 49.5, 49.8, 50.7, 51.2, 51.2, 51.7, 51.9, 51.9, 52.0, 52.1, 66.4, 66.7, 69.3, 72.0, 126.5, 126.6, 127.2, 127.7, 127.8, 128.6, 128.6, 129.1, 134.0, 134.1, 136.7, 173.8, 173.1, 173.3, 173.8.

MS (EI): *m/z* = 343 (M<sup>+</sup>, 34%), 284 (36), 253 (15), 252 (100), 199 (66), 198 (36), 192 (12), 157 (13), 156 (13), 122 (13), 115 (18).

HRMS (DIP) calcd. for C<sub>20</sub>H<sub>25</sub>NO<sub>4</sub>: 343.1784; found: 343.1785.

**Diisobutyl (1S\*,2S\*,3S\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (endo-18) and diisobutyl (1R\*,2R\*,3R\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (exo-18)**

Yield: 49 mg (37%), yellow oil.

IR (neat): 2961, 2935, 1731, 1469, 1379, 1169, 1002, 967, 748, 693 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  (mixture of *endo:exo*' 1:0.5, difficult assignment) = 0.79 (s, 1H), 0.80 (s, 1H), 0.82 (s, 1H), 0.83 (s, 1H), 0.90 (s, 3H), 0.92 (s, 3H), 0.94 (s, 3H), 0.96 (s, 3H), 1.07–1.15 (m, 1H), 1.16–1.29 (m, 2H), 1.41–1.31 (m, 1H), 1.48 (ddt,  $J$  = 9.2, 6.3, 3.6 Hz, 2H), 1.69 (dd,  $J$  = 12.1, 3.3 Hz, 0H), 1.73–1.87 (m, 1H), 1.89–2.03 (m, 2H), 2.36–2.53 (m, 1H), 2.78–2.96 (m, 2H), 3.07 (t,  $J$  = 7.7 Hz, 1H), 3.16 (ddd,  $J$  = 11.3, 8.8, 2.8 Hz, 1H), 3.27 (dd,  $J$  = 7.4, 4.3 Hz, 1H), 3.62–3.81 (m, 1H), 3.82–3.99 (m, 4H), 4.03–4.19 (m, 1H), 6.09 (dd,  $J$  = 15.7, 9.8 Hz, 1H), 6.30 (dd,  $J$  = 15.7, 9.5 Hz, 1H), 6.52 (d,  $J$  = 15.6 Hz, 1H), 6.57 (d,  $J$  = 15.7 Hz, 1H), 7.16–7.47 (m, 8H).

<sup>13</sup>C-NMR:  $\delta$  (mixture of *endo:exo*' 1:0.5, difficult assignment) = 19.1, 19.2, 19.3, 23.9, 24.0, 24.4, 27.3, 27.7, 27.8, 27.9, 30.6, 47.7, 47.9, 48.2, 49.4, 51.8, 52.7, 61.3, 63.4, 66.2, 67.2, 71.1, 71.2, 125.1, 126.6, 127.7, 127.8, 128.6, 128.8, 129.2, 133.1, 134.6, 136.6, 136.7, 171.9, 172.8, 173.3.

MS (EI):  $m/z$  = 427 (M<sup>+</sup>, 27%), 354 (15), 337 (22), 336 (100), 326 (39), 324 (13), 224 (22), 199 (38), 198 (21), 122 (15).

HRMS (DIP) calcd. for C<sub>26</sub>H<sub>37</sub>NO<sub>4</sub>: 427.2723; found: 427.2720.

**Diisobutyl (1S\*,2S\*,3R\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (endo-18) and diisobutyl (1R\*,2R\*,3S\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-1,2-dicarboxylate (exo-18)**

Yield: 50 mg (38%), yellow oil.

IR (neat): 2960, 1729, 1469, 1383, 1168, 1002, 968, 738, 692 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  (mixture of *endo:exo* 0.65:1 difficult assignment) = 0.74 (s, 1H), 0.75 (s, 1H), 0.76 (s, 1H), 0.77 (s, 1H), 0.85 (d,  $J$  = 0.8 Hz, 2H), 0.87 (d,  $J$  = 0.8 Hz, 2H), 0.92 (d,  $J$  = 0.6 Hz, 3H), 0.94 (d,  $J$  = 0.6 Hz, 3H), 0.95 (s, 2H), 0.97 (s, 2H), 1.12–1.31 (m, 2H), 1.37–1.67 (m, 3H), 1.71–2.03 (m, 7H), 2.04–2.11 (m, 1H), 2.20 (td,  $J$  = 10.4, 2.5 Hz, 1H), 2.43 (ddd,  $J$  = 10.7, 8.2, 2.4 Hz, 1H), 2.92–3.07 (m, 1H), 3.11 (d,  $J$  = 10.9 Hz, 1H), 3.18–3.48 (m, 4H), 3.69–3.82 (m, 2H), 3.84–3.95 (m, 4H), 5.97 (dd,  $J$  = 15.8, 8.7 Hz, 1H), 6.22 (dd,  $J$  = 15.8, 8.5 Hz, 1H), 6.55 (d,  $J$  = 15.8 Hz, 1H), 6.56 (d,  $J$  = 15.8 Hz, 1H), 7.20–7.43 (m, 8H).

<sup>13</sup>C-NMR:  $\delta$  (mixture of *endo:exo* 0.65:1 difficult assignment) = 19.1, 19.2, 19.3, 24.1, 24.4, 24.8, 25.0, 27.6, 27.7, 27.8, 28.6, 30.6, 49.5, 49.9, 50.9, 51.2, 51.4, 51.7, 66.5, 66.6, 69.3, 71.0, 71.1, 71.2, 72.0, 126.6, 127.4, 127.7, 128.6, 129.6, 133.9, 134.1, 136.7, 136.8, 172.5, 172.9, 173.5.

MS (EI):  $m/z$  = 427 (M<sup>+</sup>, 34%), 354 (23), 337 (22), 336 (100), 326 (47), 252 (17), 224 (24), 199 (80), 198 (35), 122 (17).

HRMS (DIP) calcd. for C<sub>26</sub>H<sub>37</sub>NO<sub>4</sub>: 427.2723; found: 427.2720.

**tert-Butyl (2S\*,3S\*,8aR\*)-3-[(E)-styryl]octahydroindolizine-2-carboxylate (endo-19)**

Yield: 20 mg (19%), yellow sticky oil.

IR (neat): 2931, 1723, 1366, 1148, 966 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.23–1.39 (m, 2H, NCHCH<sub>2</sub>CH<sub>2</sub>), 1.43 (s, 9H, *t*-Bu), 1.47–1.78 (m, 5H, CH<sub>2</sub>CHCO<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.28 (ddd,  $J$  = 12.5, 10.3, 6.3 Hz, 1H, CH<sub>2</sub>CHCO<sub>2</sub>), 2.58 (td,  $J$  = 12.3, 3.3 Hz, 1H, NCH<sub>2</sub>), 2.72 (ddd,  $J$  = 10.3, 7.4, 5.9 Hz, 1H, CHCO<sub>2</sub>), 2.86–3.01 (m, 2H, NCHCH<sub>2</sub>, NCH<sub>2</sub>), 3.98 (dd,  $J$  = 9.2, 5.9 Hz, 1H, NCH), 6.17 (dd,  $J$  = 15.7, 9.2 Hz, 1H, PhCHCH), 6.54 (d,  $J$  = 15.7 Hz, 1H, PhCH), 7.18–7.43 (m, 5H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 22.4 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.4 (NCH<sub>2</sub>CH<sub>2</sub>), 28.3 (CH<sub>3</sub>), 29.9 (NCHCH<sub>2</sub>CH<sub>2</sub>), 34.7 (CH<sub>2</sub>CHCO<sub>2</sub>*t*-Bu), 46.8 (NCH<sub>2</sub>), 49.6 (CHCO<sub>2</sub>*t*-Bu), 59.8 (NCH), 66.0 (NCH), 80.6 (CMe<sub>3</sub>), 126.5, 127.6, 128.7, 130.3, 132.7, 137.0 (ArC, C=C), 174.0 (CO).

MS (EI):  $m/z$  = 327 (M<sup>+</sup>, 18%), 271 (27), 270 (100), 254 (16), 226 (18), 180 (74).

HRMS (DIP) calcd. for C<sub>21</sub>H<sub>29</sub>NO<sub>2</sub>: 327.2198; found: 327.2199.

**(1S\*,2R\*,3S\*,8aR\*)-1-Nitro-2-phenyl-3-[(E)-styryl]octahydroindolizine (endo-20)**

Yield: 19 mg (18%), brown sticky oil.

IR (neat): 2938, 2855, 1717, 1549, 1496, 1449, 1362, 1264, 1144, 967, 736, 694 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.25–1.36 (m, 2H, NCHCH<sub>2</sub>), 1.51–1.62 (m, 1H, NCHCHCH<sub>2</sub>), 1.70–1.78 (m, 1H, NCHCHCH<sub>2</sub>), 1.84–1.93 (m, 2H, NCHCHCH<sub>2</sub>), 2.44 (ddd,  $J$  = 11.9, 8.9, 6.5 Hz, 1H, NCH<sub>2</sub>), 2.90–3.03 (m, 1H, NCH<sub>2</sub>), 3.30–3.42 (m, 1H, NCHCH<sub>2</sub>), 4.25 (dd,  $J$  = 10.1, 7.8 Hz, 1H, NCHCHPh), 4.59 (dd,  $J$  = 7.8, 7.2 Hz, 1H, NCHCHPh), 5.54 (dd,  $J$  = 8.4, 7.2 Hz, 1H, CHNO<sub>2</sub>), 5.89 (dd,  $J$  = 15.6, 10.1 Hz, 1H, PhCHCH), 6.32 (d,  $J$  = 15.6 Hz, 1H, PhCH), 7.06–7.38 (m, 10H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 23.8 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 23.9 (NCH<sub>2</sub>CH<sub>2</sub>), 26.0 (NCHCH<sub>2</sub>), 48.4 (NCH<sub>2</sub>), 51.7 (NCHCH), 62.3 (NCH), 69.2 (NCH), 93.0 (CNO<sub>2</sub>), 124.8, 126.5, 127.4, 127.8, 128.6, 128.7, 128.8, 135.2, 136.6, 136.8 (ArC, C=C).

MS (EI):  $m/z$  = 348 (M<sup>+</sup>, 1%), 303 (25), 302 (100), 300 (11), 257 (10), 219 (15), 143 (11), 117 (20), 115 (21).

HRMS (DIP) calcd. for C<sub>22</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>: 348.1838; found: 348.1825.

**(3aS\*,4R\*,9aR\*,9bR\*)-2,4-Diphenyloctahydro-1H-pyrrolo[3,4-*a*]indolizine-1,3(2H)-dione (endo-21)**Yield: 58 mg (54%), white solid, mp 151–154 °C (Et<sub>2</sub>O).IR (neat): 2943, 2850, 1701, 1497, 1393, 1189, 848, 755, 693 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 0.89–1.12 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.11–1.29 (m, 1H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.33–1.56 (m, 2H, NCHCH<sub>2</sub>), 1.60–1.84 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>), 1.97–2.20 (m, 1H, NCH<sub>2</sub>), 2.87 (ddt,  $J$  = 14.5, 6.4, 2.7 Hz, 2H, NH<sub>2</sub>, NCHCH<sub>2</sub>), 3.35–3.73 (m, 2H, PhCHCH, CH<sub>2</sub>CHCH), 4.69 (d,  $J$  = 0.9 Hz, 1H, NCHPh), 7.07–7.19 (m, 2H, ArH), 7.30–7.54 (m, 8H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 24.2 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 24.9 (NCH<sub>2</sub>CH<sub>2</sub>), 29.1 (NCHCH<sub>2</sub>), 48.5 (NCH<sub>2</sub>), 48.7 (NCHCHCO), 50.3 (NCHCHCO), 59.9 (NCH), 69.4 (NCH), 126.7, 128.0, 128.4, 128.6, 128.6, 128.7, 132.3, 136.6 (ArC), 176.1, 178.1 (2xNCO).

MS (EI):  $m/z$  = 346 (M<sup>+</sup>, 61%), 345 (48), 269 (22), 198 (12), 173 (57), 172 (100), 115 (14).

HRMS (DIP) calcd. for C<sub>22</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub>: 346.1681; found: 346.1668.

**(3aR\*,4R\*,9aR\*,9bS\*)-2,4-Diphenyloctahydro-1H-pyrrolo[3,4-*a*]indolizine-1,3(2H)-dione (exo-21)**

Yield: 26 mg (24%), yellow oil.

IR (neat): 2934, 2854, 1712, 1496, 1376, 1173, 734, 698 cm<sup>-1</sup>.

<sup>1</sup>H-NMR:  $\delta$  = 1.29–1.39 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.58–1.64 (m, 1H, NCHCH<sub>2</sub>), 1.82–1.97 (m, 2H, NCH<sub>2</sub>CH<sub>2</sub>), 2.22–2.41 (m, 2H, NCHCH<sub>2</sub>, NH<sub>2</sub>), 2.84 (d,  $J$  = 11 Hz, 1H, NH<sub>2</sub>), 3.35–3.73 (m, 1H, NCHCH), 3.17 (dd,  $J$  = 9.3, 8.5 Hz, 1H, CH<sub>2</sub>CHCH), 3.27 (dd,  $J$  = 9.3, 6.9 Hz, 1H, PhCHCH), 3.49 (d,  $J$  = 6.9 Hz, 1H, NCHPh), 7.28–7.42 (m, 6H, ArH), 7.44–7.53 (m, 4H, ArH).

<sup>13</sup>C-NMR:  $\delta$  = 24.1 (NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 25.0 (NCH<sub>2</sub>CH<sub>2</sub>), 31.3 (NCHCH<sub>2</sub>), 50.2 (NCHCHCO), 50.9 (NCH<sub>2</sub>), 53.0 (NCHCHCO), 67.8 (NCH), 72.0 (NCH), 126.6, 127.8, 127.9, 128.1, 128.7, 128.9, 129.2, 131.9 (ArC), 176.1, 176.6 (2xNCO).

MS (EI):  $m/z$  = 346 (M<sup>+</sup>, 73%), 345 (56), 269 (23), 198 (20), 173 (55), 172 (100), 115 (15).

HRMS (DIP) calcd. for C<sub>22</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub>: 346.1681; found: 346.1668.

**Acknowledgments**

The Spanish Ministerio de Ciencia e Innovación (MICINN) (projects CTQ2010-20387, and Consolider Ingenio 2010, CSD2007-00006), the Spanish Ministerio de Economía y Competitividad (MINECO) (projects CTQ2013-43446-P and CTQ2014-51912-REDC), FEDER, the Generalitat Valenciana (PROMETEO 2009/039 and PROMETEOII/2014/017), and the University of Alicante. L. C. M. thanks Spanish Government for a fellowship.

**Supporting Information**

YES (this text will be updated with links prior to publication)

**References**

**Checklist (have these on hand for manuscript submission in ScholarOne):**

- cover letter, including a statement of the work's significance
- full mailing address, telephone and fax numbers, and e-mail address of the corresponding author
- email address for each author
- original Word file
- original graphics files zipped into one zip file
- eye-catching graphical abstract as an individual file
- 5–8 key words
- separate Supporting Information file
- separate zipped Primary Data files including cover sheet (optional)

**Useful links:**

- [SYNTHESIS homepage](#)
- [SYNTHESIS information and tools for authors](#)
- [Graphical abstract samples](#) (PDF file download)
- [What is "Primary Data"?](#)
- [ScholarOne](#) (manuscript submission)

<sup>1</sup> Michael, J. P. *J. Nat. Prod.*, **2008**, *25*, 139-165.

<sup>2</sup> (a) Pansare, S. V.; Thorat, R. G. *Targets in Heterocyclic Systems* **2013**, *17*, 57. (b) Bhat, C.; Tilve, S. G. *RSC Adv.*, **2014**, *4*, 5405. (c) Bronner, S. M.; Im, G.-Y. J.; Garg, N. K. *Heterocycles in Natural Product Synthesis* **2011**, 221. (d) Gerber-Lemaire, S.; Juillerat-Jeanneret, L. *Chimia* **2010**, *64*, 634-639.

<sup>3</sup> Procedures employing an intramolecular cyclization affording simultaneously 5 and 6 membered rings have seldom been reported. See, for example: Brambilla, M.; Davies, S. G.; Fletcher, A. M.; Roberts, P. M.; Thomson, J. E. *Tetrahedron* **2014**, *70*, 204-211.

<sup>4</sup> Bhat, C.; Tilve, S. G. *Chem. Commun.* **2014**, *4*, 5405-5452.

<sup>5</sup> See, for example: (a) Pronin, S. V.; Tabor, M. G.; Jansen, D. J.; Shenvi, R. A. *J. Am. Chem. Soc.* **2012**, *134*, 2012. (b) Shapland, P. *Nature Chem.* **2012**, *4*, 441-442.

<sup>6</sup> See, for example: (a) Vu, H.-D.; Renault, J.; Roisnel, T.; Gouault, N.; Uriac, P. *Tetrahedron Lett.* **2016**, *57*, 7, 3036-3038. (b) Tan, Y.; Chen, Y.-J.; Lin, H.; Sun, X.-W.; Yang, X. D.; Lin, G.-Q. *Chem. Commun.* **2014**, *50*, 15913-15915.

<sup>7</sup> For general reviews dealing with general 1,3-DC, see (a) Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry towards Heterocycles and Natural Products; Padwa, A., Pearson, W. H., Eds.; John Wiley & Sons: New Jersey, 2003; (b) Nájera, C.; Sansano, J. M. *Curr. Org. Chem.* **2003**, *7*, 1105-1150; (c) Eberbach, W. *Sci. Synth.* **2004**, *27*, chp. 11, 441. (d) Coldham, I.; Hufton, R. *Chem. Rev.* **2005**, *105*, 2765-2810; (e) Nair, V.; Suja, T. D. *Tetrahedron* **2007**, *63*, 12247-12275; (f) Padwa, A.; Bur, S. K. *Tetrahedron* **2007**, *63*, 5341-5378.

<sup>8</sup> (a) Pearson, W. H. *Pure Appl. Chem.* **2002**, *74*, 1339-1347. (b) Pearson, W. H.; Stoy, P. *Synlett* **2003**, 903-921.

<sup>9</sup> For natural product synthesis see, for example: Shanahan, C. S.; Fang, C.; Paull, D. H.; Martin, S. F. *Tetrahedron* **2003**, *69*, 7592-7607.

<sup>10</sup> For an example dealing with a 5→6 sequence using 1,3-DC of azomethine ylides, see: Koley, D.; Srinivas, K.; Krishna, Y.; Gupta, A. *RSC Adv.* **2004**, *4*, 3934-3937.

<sup>11</sup> Coldham, I.; Jana, S.; Watson, L.; Pilgram, C. D. *Tetrahedron Lett.* **2008**, *49*, 5408-5410.



- <sup>12</sup> (a) Grigg, R.; Surendrakumar, S.; Thianpatanagul, S.; Vipond, D. *J. Chem. Soc., Perkin Trans. 1* **1988**, 2693-2701. (b) Grigg, R.; Surendrakumar, S.; Thianpatanagul, S.; Vipond, D. *J. Chem. Soc., Chem. Commun.* **1987**, 47-49.
- <sup>13</sup> The multicomponent reaction from cyclic secondary  $\alpha$ -amino esters and bifunctional carbonyl compounds occurred through 1,5-H shifts followed by reaction with dipolarohiles: Grigg, R.; Rankovic, Z.; Thornton-Pett, M.; Somasunderam A. *Tetrahedron* **1992**, *48*, 10431-10442.
- <sup>14</sup> Combinations of imine-azomethine ylide-cycloaddition cascades with acid catalysed Pictet-Spengler spirocyclisation affording novel polycyclic spirocycles has been reported: Grigg, R.; Thornton-Pett, M.; Yoganathan, G. *Tetrahedron* **1999**, *55*, 8129-8140.
- <sup>15</sup> Dondas, H. A.; Duraisingham, J.; Grigg, R.; MacLachlan, W. S.; MacPherson, D. T.; Thornton-Pett, M.; Sridharan, V.; Suganthan, S. *Tetrahedron* **2000**, *56*, 4063-4070.
- <sup>16</sup> Joucla, M.; Mortier, J.; Hamelin, J. *Tetrahedron Lett.* **1985**, *26*, 2775-2778
- <sup>17</sup> (a) Mancebo-Aracil, J.; Nájera, C.; Sansano, J. M. *Chem. Commun.* **2013**, *49*, 11218. (b) Sengupta, T.; Khamarui, S.; Samanta, S.; Maiti, D. K. *Chem. Commun.* **2013**, *49*, 9962. (c) Mancebo-Aracil, J.; Nájera, C.; Castelló, L. M.; Sansano, J. M.; Larrañaga, O.; de Cózar, A.; Cossío, F. P. *Tetrahedron* **2015**, *71*, 9645-9661.
- <sup>18</sup> Chalet, Ch.; Croix, C.; Alagille, D.; Normand, S.; Delwail, A.; Favot, L.; Lecron, J.-C.; Viaud-Massuard, M.-C. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 1019-1022.
- <sup>19</sup> This *exo*-diastereoselectivity has been observed in the synthesis of pyrrolizidines with chalcone. Biswajit Gayen, Avijit Banerji & Kaliprasanna Dhara, *Synth. Commun.* **2016**, DOI:10.1080/00397911.2015.1135954.
- <sup>20</sup> Available structure deposited in CCDC with number 1496416.
- <sup>21</sup> For an analogous approach using decarboxylation of proline derivatives see for example: Kang, T.-R; Cheng, Y.; He, L.; Ye, J.; Liu, Q.-Z. *Tetrahedron Lett.* **2012**, *53*, 2552-2555 and references cited therein.